

Sponsor	1100021	Albemarle Corporation	Create Date:	4/6/01
CAS Number	2194853	Cyclododecane, 1,2,5,6,9,10-hexabromo-		
Consortium	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)		

Select an End Point: TO - Genetic toxicity in vivo

Check for Conflicts

Physical/Chemical Properties <input checked="" type="checkbox"/> Melting Point <input checked="" type="checkbox"/> Partition Coefficient <input checked="" type="checkbox"/> Water Solubility <input type="checkbox"/> Boiling Point <input checked="" type="checkbox"/> Vapor Pressure	Ecotoxicity <input checked="" type="checkbox"/> Acute Toxicity to Fish <input checked="" type="checkbox"/> Acute Toxicity to Aquatic Invertebrates <input checked="" type="checkbox"/> Toxicity to Aquatic Plant
Environmental Fate <input type="checkbox"/> Photodegradation <input checked="" type="checkbox"/> Biodegradation <input type="checkbox"/> Stability in Water <input checked="" type="checkbox"/> Transport	Health <input checked="" type="checkbox"/> Acute Toxicity <input checked="" type="checkbox"/> Repeat Dose Toxicity <input type="checkbox"/> Reproductive Toxicity <input checked="" type="checkbox"/> Developmental Tox/Teratogenicity <input checked="" type="checkbox"/> Genetic Toxicity in Vivo <input checked="" type="checkbox"/> Genetic Toxicity in Vitro

02 JAN -6 PM 12:41

RECEIVED
APPT NCIC

A12201-13459B

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Melting Point

Sponsor ID 110002

Albemarle Corporation

Create Date 4/6/01

CAS Number 3194556

Cyclododecane, 1,2,5,6,9,10-hexabromo-

Study Number 1

Consortia ID 1101012

CMA Brominated Flame Retardant Industry Panel (BFRIP)

Completed: Y

Revision Date:

12/5/01

Test Substance

Remarks

The test substance consisted of various commercial products.

Chemical Category

Method

>> Method/Guideline followed

Not specified.

>> GLP Unknown

>> Year study performed 1994

Remarks for Method

Results

>> Precision range

>> Melting Point Value 175

>> Upper Value 195

>> Unit °C

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Melting Point

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/12/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	Y

>> Decomposition Yes

>> Sublimation No

Results Remark

Various melting points have been reported for different products: 175-183 degrees C (Saytex HBCD-LM); 187-195 degrees C (Saytex-HM), 190 degrees C (GLCC product).

Conclusions

HBCD is a solid at room temperature whose melting point varies with composition.

Data Quality

Reliability Good

Data Reliability Remarks

The melting point data was provided by commercial manufacturers of the substance.

Reference

>> Remarks

IUCLID Dataset. Substance ID: 25637-99-4. 18-Feb-2000.

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Melting Point

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

General

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Partition Coefficient

Sponsor ID	1100021	Albemarle Corporation	Create Date	11/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OPPTS 830.7560 Partition Coefficient (n-Octanol/Water), Generator Column Method

>> GLP Yes

>> Year study performed 1997

Remarks for Method

A single generator column was prepared for the definitive test. The column was packed with Chromosorb W HP support and loaded with an approximate 0.2% solution of the test substance in octanol. Dilutions of the test substance solution in octanol were analyzed. The column temperature was maintained at 25 +/- 0.05 degrees C and reagent water saturated with octanol was pumped through it at approximately 1 mL per min to elute the test substance. Samples of the eluate were collected and analyzed to determine the concentration of the test substance in the aqueous fractions.

The analytical method consisted of extracting the aqueous samples with dichloromethane (DCM), evaporating the DCM, and reconstituting the sample residues in acetonitrile/water (50:50, v/v).

Results

>> Precision

=

>> Value of Log Pow

5.625

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Partition Coefficient

Sponsor ID	1100021	Albemarle Corporation	Create Date	11/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	Y

>> Upper Value 0

>> Temperature 25 degrees C

Results Remark

HBDC's water solubility was previously determined to be 0.0034 mg/L (Stenzel and Markley, 1997).

No interferences were observed at or above the limit of quantitation in the matrix blank sample. The percent recovery of the 1.00 and 10.0 ug HBDC/L matrix fortifications were 104 and 85%. The mean recovery was calculated at 95% of nominal.

The nominal flow rate of reagent water through the generator column was measured prior to the start of sample collection. Flow rates were also calculated based on the volume and collection time of each sample that was analyzed. The pump setting was 1.0 mL/min and the flow rate was measured at 1.0 mL/min. The calculated flow rates for samples averaged 0.87 mL/min and ranged from 0.86 to 0.87 mL/min.

The mean concentration of HBDC measured in the aqueous samples eluted from the generator column was 3.97 ug HBDC/L or 6.19×10^{-9} M (molecular weight of HBDC is 641.7 g/mole).

The mean concentration of HBDC measured in the octanol stock solution samples was 1.67 g HBDC/L or 2.61×10^{-3} M (molecular weight of HBDC is 641.7 g/mole).

Conclusions

The octanol/water partition (Kow) coefficient was calculated from the following equation:

$$Kow = \frac{\text{Measured Concentration in Octanol (M)}}{\text{Measured Concentration in Aqueous Samples (M)}}$$

Based on the results from octanol samples collected from the stock solution and aqueous samples collected from the generator column, the mean octanol/water partition coefficient (Kow) for HBDC was determined to be 4.22×10^{-5} (log Kow = 5.625).

Data Quality

Reliability High

Data Reliability Remarks

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Partition Coefficient

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

This study was performed according to current guidelines and Good Laboratory Practices by a laboratory with considerable experience with these studies. Extensive attention was paid to analytical method development and performance.

Reference

>> Remarks

MacGregor, J and Nixon, W. (1997) Hexabromocyclododecane (HBCD): Determination of n-Octanol/Water Partition Coefficient. Project Number: 439C-104. Wildlife International LTD, Easton, MD.

General

Study sponsored by the Chemical Manufacturers Association Brominated Flame Retardant Industry Panel, Arlington, VA.

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Vapor Pressure

Sponsor ID	1100021	Albemarle Corporation	Create Date	12/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	Y

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OECD Method 104; U.S. EPA OPPTS 830.7950 Vapor Pressure

>> GLP Yes

>> Year study performed

1997

Remarks for Method

The objective of this study was to determine the vapor pressure of HBCD at ambient temperature using a spinning rotor gauge (SRG). The SRG method was chosen due to the extremely low vapor pressure anticipated for this substance.

The SRG system was configured with an empty 10 mL beaker in the sample chamber to make control measurements. The system baseline pressure and out-gassing rate were each measured twice.

A sample of the test substance in a 10 mL beaker was placed in the sample chamber. The SRG system was used to monitor the steady-state pressure of the sample while the system was open to the vacuum pumps, and the pressure increase from the sample while the valve was closed to the pumps. The steady-state pressure and pressure increase measurements were repeated three times.

Results

>> Precision

=

>> Vapor Pressure Value

0.0000627

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Vapor Pressure

Sponsor ID	1100021	Albemarle Corporation	Create Date	11/6/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

>> Upper Value 0

>> Unit Pascals

>> Temperature 21 degrees C

>> Decomposition No

Results Remark

The baseline pressure of the system containing an empty beaker was determined to be less than 1×10^{-7} Pa for both measurements. The technical specifications of the SRG indicated the low end of the measurement range to be 1×10^{-5} Pa. The baseline pressure was considered to be essentially zero, and indicated the system was free of contamination. The out-gassing rate (slope) was $<1 \times 10^{-7}$ Pa/sec for both measurements. The out-gassing rate indicated there were no leaks in the system.

The mean steady-state pressure of the HBCD sample was 6.166×10^{-5} based on three separate determinations. The slope of the line fit to the pressure increase data was less than the out-gassing rate of the empty system for each determination, indicating the system had achieved saturation of the gas from the HBCD sample and was leak-free. The intercept was only slightly greater than the steady-state pressure. The temperature of the system averaged 21 degrees C.

The vapor pressure for each determination of the HBCD sample was calculated from the following equation:

$$\text{Vapor Pressure} = \text{intercept of sample} - \text{mean intercept of empty system.}$$

The mean vapor pressure of HBCD was determined to be 6.27×10^{-5} Pa with a standard deviation of 0.21×10^{-5} .

The vapor pressures of di(2-ethyl-hexyl)phthalate and hexachlorobenzene were measured using the same SRG system and determined to be 4.3×10^{-5} Pa and 1.6×10^{-3} Pa, respectively. Both of these measurements were consistent with ranges found in the literature.

Conclusions

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Vapor Pressure

Sponsor ID	110002	Albemarle Corporation	Create Date	12/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Based on the results from three sets of measurements collected from the spinning rotor gauge, the vapor pressure of HBCD was determined to be 6.27×10^{-5} Pa at 21 degrees C.

Data Quality

Reliability High

Data Reliability Remarks

This study was performed according to current guidelines and Good Laboratory Practices by a laboratory with considerable experience with these studies. Extensive attention was paid to analytical method development and performance.

Reference

>> Remarks

Stenzel, J and Nixon, W. (1997) Hexabromocyclododecane (HBCD): Determination of the Vapor Pressure Using a Spinning Rotor Guage. Project Number: 439C-117. Wildlife International LTD, Easton, MD.

General

Study sponsored by the Chemical Manufacturers Association Brominated Flame Retardant Industry Panel.

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Water Solubility

Sponsor ID	1100021	Albemarle Corporation	Create Date	10/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	N

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OECD Method 105, U.S. EPA 40 CFR Ch. 1 Section 796.1860 Water Solubility- Generator Column Method

>> GLP Yes

>> Year study performed

1997

Remarks for Method

This study was performed according to OECD Method 105 and U.S. EPA 40 CFR Ch. 1 Section 796.1860 Water Solubility- Generator Column Method.

A generator column was prepared. The column temperature was maintained at 25.0 degrees C and reagent water was pumped through it at approximately 2 mL per minute to elute the test substance. Samples of the eluate were collected and analyzed to determine the saturation concentration of the test substance. The flow rate of reagent water through the column was reduced to approximately half the original flow rate and the saturation concentration determined again.

Results

>> Precision

<

>> Water Solubility Value

1

>> Upper Value

0

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Water Solubility

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

>> Unit mg/L

>> Temperature 25 degrees C

>> Solubility Category Insoluble

>> pH Value 8

>> pKa Value 0

Results Remark

No interferences were observed at or above the limit of detection (0.5 ug HBCD/L) in any of the matrix blank or reagent blank samples. The peak area response for the matrix blanks was always below the response of the lowest calibration standard. The mean recovery from 10 matrix samples fortified at 10 ug/L was 105% (standard deviation 2.0), and ranged from 103% to 108%. The mean recovery from 10 matrix samples fortified at 1 ug/L was 104% (standard deviation 5.2), and ranged from 100% to 110%. The 1 ug/L concentration was considered the limit of quantitation.

A brief description of the analytical method is as follows: samples were extracted using dichloromethane (DCM). The DCM was evaporated to dryness and 1.0 ml of acetonitrile/water (50:50) was added. The samples were analyzed using HPLC/UV.

The nominal flow rate of reagent water through the generator column was initially set at 1.0 mL/min. The initial flow rate was measured at 2.0 mL/min prior to the start of sample collection. Samples were collected at this flow rate until the solubility plateau was achieved. The calculated flow rates for samples collected at the initial flow rate averaged 1.96 mL/min (range 1.88-1.98 mL/min). After the solubility plateau was achieved, the flow rate was reduced to ~half the initial flow rate. The reduced flow rate was measured at 1.0 mL/min prior to resuming sample collection. The calculated flow rates averaged 0.92 mL/min (range 0.91-0.93).

All samples collected at a nominal flow rate of 2.0 mL/min were analyzed and the solubility limit was considered to have been achieved when at least 5 consecutive samples gave similar results. The mean concentration in samples meeting this criteria was 0.0034 mg/L with a standard deviation of 0.23.

The results from analyses of samples eluted at a nominal flow rate of 1.0 mL/min found a mean

EPA High Production Volume (HPV) Track

Physical-Chemical End Point:
Water Solubility

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

HBCD concentration of 0.0033 mg/L with a standard deviation of 0.20.

The pH of the water obtained from Wildlife International Ltd's well in February/March 1997 had a mean pH of 8.3 (range 8.2-8.4).

NOTE: HBCD has no ionizable groups and therefore the pKa value does not apply. A value of 0 was entered in the "pka value" field because this was a mandatory numeric field.

Conclusions

The solubility of HBCD in water was determined to be 0.0034 +/- 0.2 mg/L at 25 degrees C.

Data Quality

Reliability High

Data Reliability Remarks

This study was performed according to current guidelines and Good Laboratory Practices by a laboratory with considerable experience with these studies. Extensive attention was paid to analytical method development and performance.

Reference

>> Remarks

Stenzel, J. And Markley, B. (1997) Hexabromocyclododecane (HBCD): Determination of the Water Solubility. Project Number: 439C-105. Wildlife International LTD, Easton, MD.

General

Study sponsored by the Chemical Manufacturer's Association Brominated Flame Retardant Industry Panel, Arlington, VA.

EPA High Production Volume (HPV) Track

Environmental Fate and Pathway End Point:
Biodegradation

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	319156	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

EPA OPPTS Method 835.3200: Ready Biodegradability, Closed Bottle Test; OECD Guideline 301D

>> Test Type

aerobic

>> GLP Yes

>> Year study performed 1996

>> Contact Time 28

>> Inoculum

activated sludge, domestic, adapted

Remarks for Method

The test contained an inoculum control group, a reference group and a treatment group. The blank control, reference, and treatment groups contained ten replicate test chambers. The inoculum control was used to measure the dissolved oxygen consumption of the inoculum and was not dosed with a carbon source. The reference chambers were dosed with sodium benzoate, a substance known to be biodegradable, at a concentration of 2 mg/L. The treatment group test chambers were used to evaluate the test substance at 7.7 mg/L. Measurements of oxygen consumption were performed on two test chambers from the control, reference and treatment groups on days 0, 7, 14, 21, and 28.

The test inoculum was secondary clarifier supernatant collected from Prospect Bay Wastewater Treatment Facility, Grasonville, MD. The theoretical oxygen demand value used to calculate the percent degradation of the test substance was 0.75 mg O₂/mg.

Results

12/20/01

Page 1 of 3

EPA High Production Volume (HPV) Track

Environmental Fate and Pathway End Point:
Biodegradation

Spenser ID	1100021	Albemarle Corporation	Create Date	11/6/01
CAS Number	3191356	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

>> Precision =

>> Degradation Value 0

>> Upper value 0

>> Time Frame 28

>> Time Units Days

>> Breakdown products No

Results Remarks

The temperature range recorded during the test was 18-20 degrees C. The result of the standard plate count performed on the inoculum was 3.7×10^4 CFU/ml.

The average oxygen uptake exhibited by the control, reference, and treatment groups was measured at 0, 7, 14, 21 and 28 days. The oxygen depletion of the inoculum control was less than or equal to 1.5 mg O₂/L. Degradation of the test substance was not observed over the 28-day test period.

The viability of the inoculum and validity of the test was supported by the results of the reference substance, sodium benzoate, degrading approximately 94%. An average percent biodegradation of > 60% was achieved by day 7, thereby fulfilling the criteria for a valid test.

Conclusions

Degradation of the test substance, HBCD, at 7.7 mg/L was not observed over the 28-day test period.

EPA High Production Volume (HPV) Track

Environmental Fate and Pathway End Point:
Biodegradation

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	319356	Cyclododecane 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Data Quality

Reliability High

Data Reliability Remarks

This study was performed according to current guidelines and Good Laboratory Practices by a laboratory with considerable experience with these studies. Extensive attention was paid to analytical method development and performance.

Reference

>> Remarks

Schaefer, E and Haberlein, D. (1996) Hexabromocyclododecane (HBCD): Closed Bottle Test. Project No.: 439E-102. Wildlife International Ltd. Easton, MD.

General

Study sponsored by the Chemical Manufacturers Association Brominated Flame Retardant Industry Panel.

EPA High Production Volume (HPV) Track

Environmental Fate and Pathway End Point: Transport
between Environmental Compartments (Fugacity)

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

12/18/01

Test Substance

Remarks

Hexabromocyclododecane (HBCD)

Chemical Category

Method

>> Method/Guideline followed

Developed by D. Mackay and co-workers

>> Test Type

Level III fugacity model

>> Year study performed

2001

Remarks for Method

Model Used: Level III Fugacity Model (Full-Output), EPIWIN V3.04

Input parameters: chemical structure only; model default parameters accepted; model based on emissions of 1000 kg/hr each to air, water and soil.

Results

>> Media

Air: 0.000685%; Water: 2.06%; Soil: 40.1%; Sediment: 57.9%

>> Distribution Concentration

Not provided by model.

EPA High Production Volume (HPV) Track

Environmental Fate and Pathway End Point: Transport between Environmental Compartments (Fugacity)

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Results Remark

Estimated by model:
Soil Koc: $2.25 \times 10^+7$
Vapor Pressure: 1.68×10^{-8} mmHg
Liquid VP: 5.74×10^{-7} mm Hg (super-cooled)
Melting Pt: 180 deg C
Log Kow: 7.74
Henry's LC: 6.43×10^{-11} atm-m³/mole

Conclusions

If released at equal rates to air, water and soil, HBCD is predicted to partition primarily to sediment (appr. 58%) and soil (appr. 40%). Only appr. 2% would partition to water with only trace (appr. 0.0007%) amounts to air. Appr. 90% would be reacted with only appr. 11% advected.

The model was also run 7 times using all permutations of air, water and soil emission rates as either 0 or 1000 kg/hr. The results were as follows. If released solely to air, the model predicted HBCD would partition appr. two-thirds to soil and one-third to sediment; 97% reacted. If released solely to water, HBCD would partition to sediment; total reacted = 71%. If released solely to soil, HBCD would remain in soil; total reacted = 100%. If released at equal rates to both air and water, HBCD would partition two-thirds to sediment and one-third to soil; total reacted = 84%. If released at equal rates to both air and soil, HBCD would partition appr. 80% to soil and 12% to sediment; total reacted = 98%. If released to water and soil, HBCD would partition two-thirds to sediment and one-third to soil; total reacted = 86%.

Based on the above, HBCD is not expected to move from water, soil or sediment to air. Furthermore, HBCD is not expected to move from soil into water.

Data Quality

Reliability

High

Data Reliability Remarks

Reference

EPA High Production Volume (HPV) Track

Environmental Fate and Pathway End Point: Transport between Environmental Compartments (Fugacity)

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

>> Remarks

Level III Fugacity Model, EPIWIN V3.04, Syracuse Research Corporation, Syracuse, NY.

General

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Fish

Sponsor ID	11000121	Albemarle Corporation	Create Date	4/6/01
CAS Number	1193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	N

Revision Date

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OECD Method 203

>> Test Type

flow-through

>> GLP Yes

>> Year study performed 1997

>> Species

Oncorhynchus mykiss

>> Analytical monitoring HPLC/UV/VIS Detector; LOQ=0.04 ug/l

>> Exposure period 96 hours

>> Statistical Method None needed - no mortality observed.

Remarks for Method

This study was performed according to OECD Method 203 and TSCA Title 40 of CFR, Part 797, Section 1400. Rainbow trout were exposed to one of five test concentrations, a solvent control, or the negative (well water) control. Two replicate test chambers were maintained in each treatment and control group. Ten rainbow trout were used in each test chamber for a total of 20 rainbow trout per test concentration. Nominal test concentrations were selected in consultation with the Sponsor, and were based on the solubility of the test compound in water (3.4 ug/L) and the results of an exploratory rangefinding test. Due to co-eluting artifacts at 96

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Fish

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	31915-00-0	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	N

hrs, mean measured test concentrations were determined analytically from samples of test water collected from each treatment and control group at the beginning of the test and at approximately 48 hrs.

The selection of exposure concentrations took into consideration the water solubility limit and a finding of no acute toxicity from an exploratory rangefinding test. The water solubility limit was determined in a generator column elution study to be 3.4 ug/L. However, there was a potential to have a slight enhancement of HBCD's water solubility due to the use of dimethylformamide (DMF) as a vehicle in the diluter system. For this reason, the highest test concentration selected for the acute toxicity test was twice the defined solubility limit (i.e., 6.8 ug/L). The series of 5 nominal test concentrations used in the test were 1.5, 2.2, 3.2, 4.6 and 6.8 ug/L. In this way, the solubility limit of HBCD was bracketed by the five concentrations.

Delivery of the test substance was initiated approximately 6 days prior to the introduction of the fish to the test water in order to achieve equilibrium of the test substance in the test chambers. The fish were indiscriminately assigned to exposure chambers at test initiation. Observations of mortality and other clinical signs were made approximately 1, 24, 48, 72 and 96 hrs after test initiation. The no mortality concentration and no observed effect concentration (NOEC) were determined by visual interpretation of the mortality and clinical observation data.

All fish were from the same source and year class, and the total length of the longest fish was no more than twice the length of the shortest. The average length of 10 negative control fish at the end of the test was 55 mm with a range of 50-61 mm. The wet weight of 10 negative control fish at the end of the test was 2.4 g with a range of 1.6-3.6 g. Loading, defined as the total wet weight of fish per liter of test water that passed through the test chamber in 24 hrs, was 0.27 g fish/L/day.

Temperature, dissolved oxygen, and pH were measured. Temperatures were within the limits of the 12 +/- 2 degrees C range established for the test. Dissolved oxygen concentrations were greater than or = 78% of saturation throughout the test. Water pH ranged from 8.2-8.3. Total organic carbon values were <1.0 mg C/L at test initiation and termination.

Test substance concentrations were determined via HPLC using a UV/VIS detector.

Results

>> Nominal concentration 0, 0.0015, 0.0022, 0.0032, 0.0046, 0.0068

>> Measured concentration 0, 0.00075, 0.0015, 0.0023, 0.0023, 0.0025

>> Precision >

>> Endpoint Type LC0

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Fish

Sponsor ID	110002	Albemarle Corporation	Create Date	4/6/01
CAS Number	319455-1	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

>> Endpoint Value >> Unit used

>> Concentration Type >> Endpoint Time

>> Statistical results

None needed - no mortality observed.

Results Remark

One set of pretest water samples was collected from the highest and lowest test concentrations and analyzed for HBCD concentrations. All pretest samples yielded concentrations that were considerably lower than the expected concentrations. The toxicity test was initiated and measurements of the HBCD concentrations in all test chambers were made at the beginning, middle and end of the test. In general, concentrations of HBCD made on samples collected at Day 0 and Day 2 were variable and failed to correspond to the dilution series expected from the nominal concentrations. All diluter operational records were checked and no evidence of any malfunctions or errors were found. Concentrations measured in the Day 4 samples were artificially high due to co-eluting artifacts at the retention time of HBCD. Attempts were made to separate the co-eluting artifacts during a reanalysis of the original Day 4 sample extracts, but the resulting chromatography showed those same interferences.

While the pattern of measured HBCD was unexpected, the results suggest that the exposure solutions were at the solubility limit of HBCD in the diluter system. The variability in the measured concentrations could have been influenced by the temperature of the exposure water (12 degrees C), the flow-through design, or the hydrophobic nature of HBCD (as evidenced by its nonpolar alkane structure and extremely low water solubility). These factors could explain both the failure of the measured values to correspond to the nominal concentrations and the variability observed in the measured concentrations. Overall, it appears that the solubility limit of HBCD, under the conditions that it was applied in this test, is within the range of 2.0 - 3.0 ug/L. The values obtained in the Day 4 samples were not reflective of the true conditions due to the co-eluting artifacts, and therefore, were not used in the study.

Temperatures were within the limits of the 12 +/- 2 degrees C range established for the test. Dissolved oxygen concentration of > or = 78% of saturation were observed throughout the test. Water pH was consistent with values for moderately-hard water and ranged from 8.2 to 8.3. Total organic carbon values were < 1.0 mg C/L at test initiation and termination.

Observations for mortality and other signs of toxicity were made daily. Rainbow trout in the negative control and solvent control groups appeared healthy and normal throughout the test. All rainbow trout in the 1.5, 2.2, 3.2, 4.6 and 6.8 ug/L (nominal) treatment groups also appeared normal throughout the test with no mortalities or overt signs of toxicity. Based on these results, the LC50 values at 24, 48, 72 and 96 hours were estimated to be >6.8 ug/L, the highest concentration tested.

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Fish

Sponsor ID	110002	Albemarle Corporation	Create Date	11/6/01
CAS Number	318156	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	110101	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	N

Conclusions

The 96-hour LC50 value for rainbow trout exposed to HBCD was >6.8 ug/L (nominal) (>2.5 ug/L mean measured concentration), the highest concentration tested and twice HBCD's water solubility (3.4 ug/L). Based on the mortality and observation data, the 96-hour no mortality concentration and the no-observed-effect-concentration were 6.8 ug/L (nominal) (2.5 ug/L mean measured concentration) and was higher than the water solubility of HBCD.

Data Quality

Reliability

High

Data Reliability Remarks

This study was performed according to current guidelines and Good Laboratory Practices by a laboratory with considerable experience with these studies. Extensive attention was paid to analytical method development and performance.

Reference

>> Remarks

Graves, W and Swigert, J. (1997) Hexabromocyclododecane (HBCD): A 96-Hour Flow-Through Acute Toxicity Test with the Rainbow Trout (*Oncorhynchus mykiss*). Project Number: 439A-101. Wildlife International LTD, Easton, MD.

General

Study sponsored by the Chemical Manufacturers Association Brominated Flame Retardant Industry Panel, Arlington, VA.

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Aquatic Invertebrates

Sponsor ID	1102021	Albemarle Corporation	Create Date	4/6/01
CAS Number	1194546	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OECD Method 202; TSCA Title 40 CFR, Part 797, Section 1300

>> Test Type

flow-through

>> GLP

Yes

>> Year study performed

1997

>> Species

Daphnia magna

>> Analytical monitoring

HPLC; Limit of Quantitation=0.4 ug/L

>> Exposure period

48 Hours

>> Statistical Method

None - no dose response pattern

Remarks for Method

Daphnids were exposed to one of five test concentrations, a solvent control or the negative (well water) control. Two replicate test chambers were maintained for each treatment and control group. Ten daphnids were used in each test chamber for a total of 20 daphnids per test concentration. Nominal test concentrations were based upon the solubility of the test substance in water (3.4 ug/L) and the results of an exploratory rangefinding toxicity test. Nominal test concentrations were 1.5, 2.2, 3.2, 4.6 and 6.8 ug/L. Mean measured test concentrations were analytically determined (HPLC with UV/VIS detector) from samples of test water collected from each treatment and control group at the beginning and end of the test.

Results

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Aquatic Invertebrates

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

Delivery of the test substance was initiated approximately 4 days prior to the introduction of the daphnids to the test water in order to achieve equilibrium of the test substance in the test chambers. Daphnids were indiscriminately assigned to exposure chambers at test initiation. Observations of mortality/immobility and other clinical signs were made approximately 2, 24 and 48 hours after test initiation. Cumulative percent mortality and immobility observed in the treatment groups were used to estimate EC50 values at 24 and 48 hours. The no mortality/immobility concentration and the no-observed-effect concentration (NOEC) were determined by visual interpretation of the mortality, immobility and clinical observation data.

Daphnid neonates used in the test were less than 24 hours old and were obtained from cultures maintained by Wildlife International Ltd, Easton, MD. Adult daphnids were cultured in water from the same source and at approximately the same temperature as that used during the test except supplemented with selenium. Daphnids in the cultures were held for 15-29 days prior to collection of the juveniles for testing. The progeny of 7 adults were used in the test. The adults were fed prior to test initiation, but neonates were not fed during the test. During the 14-day holding period preceding the test, water temperatures ranged from 20.2 to 21.4 degrees C. The pH of the water ranged from 8.0 to 8.5. Dissolved oxygen ranged from 8.2 to 9.0 mg/L.

A continuous-flow diluter was used to deliver each concentration of the test substance, a solvent control, and a negative (dilution water) control. Syringe pumps (Harvard Apparatus) were used to deliver the five test substance stock solutions and the solvent for the solvent control into mixing chambers assigned to each treatment level and the solvent control. The stock solutions were diluted with well water in the mixing chambers in order to obtain the desired test concentrations. The flow of dilution water to the mixing chambers was controlled by rotameters. Rotameters were calibrated prior to test initiation. The flow of test water from each mixing chamber was split and allowed to flow into replicate test chambers. The proportion of test water that was split into each replicate was checked prior to the test to ensure that flow rates varied by no more than +/- 10% of the mean for the two replicates.

The diluter was adjusted so that each test chamber received ~14 volume additions of test water every 24 hours. The stock solution delivery pumps were calibrated before the test, and the general operation of the diluter was checked visually at least two times daily during the test and once at the end of the test.

Test compartments were constructed from 300 mL glass beakers ~ 8 cm in diameter and 13 cm in height. The beakers were suspended in 8-L stainless steel test chambers filled with ~6.5 L of test water. Test chambers were indiscriminately positioned in a temperature-controlled water bath designed to maintain a temperature of 20+/-1 degreeC. The water bath was enclosed in a plexiglass ventilation hood. Test chambers were labeled with the project number, test concentration, and replicate.

The water used for culturing and testing was freshwater obtained from a well ~45 meters deep located on the Wildlife International Ltd. Site. The well water is characterized as moderately-hard water. The dissolved oxygen content of the water ranged from 8.8-8.9, 9.0-9.1, and 8.8-8.9 mg/L at 0, 24, and 48 hours, respectively. The pH of the water was 8.1, 8.2-8.4, and 8.2-

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Aquatic Invertebrates

Sponsor ID	100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	31915-56	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

8.3 at 0, 24 and 48 hours, respectively. The temperature of the water ranged from 19.8-19.9 and 19.9-20.0 at 0 and 48 hours, respectively. The 0-hour dilution water measurements for hardness, alkalinity and specific conductance were 132 mg/L as CaCO₃, 176 mg/L as CaCO₃ and 320 umhos/cm, respectively.

Lighting was provided by fluorescent tubes that emitted wavelengths similar to natural sunlight. A photoperiod of 16 hours of light and 8 hours of darkness was controlled with an automatic timer. Light intensity at test initiation was ~ 242 lux at the surface of the water.

>> Nominal concentration 0, 0.0015, 0.0022, 0.0032, 0.0046, 0.0068 mg/L

>> Measured concentration 0, 0.0024, 0.0018, 0.0021, 0.0023, 0.0032 mg/L

>> Precision >

>> Endpoint Type LC0

>> Endpoint Value 0

>> Unit used mg/L

>> Concentration Type Nominal

>> Endpoint Time 48

>> Statistical results

Statistics not performed due to lack of dose response.

Results Remark

The selection of exposure concentrations took into consideration the water solubility limit (3.4 ug/L) and a finding of no acute toxicity from an exploratory range finding test. However, there was a potential to have a slight enhancement of HBCD's water solubility due to the use of dimethyl formamide (DMF) as a vehicle in the diluter system. For this reason, the highest test concentration selected was twice the defined solubility limit (i.e., 6.8 ug/L). The series of nominal test concentrations bracketed the solubility limit of HBCD by five concentrations.

Two sets of pretest samples were collected from the highest and lowest test concentrations and analyzed. The Day -3 and -2 samples indicated that the test concentrations were stable, but somewhat lower than expected. Measurements of HBCD concentration in all test chambers were made at the beginning and end of the test. These measurements indicated that HBCD concentrations were generally similar across all treatment levels, and may reflect a

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Aquatic Invertebrates

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	1103556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

phenomenon in the delivery system whereby HBCD adsorbed to the physical surfaces of the diluter system. This could be due to the hydrophobic nature of HBCD as evidenced by its nonpolar alkane structure and extremely low water solubility. This characteristic could have enabled the inert surfaces (e.g. Stainless steel and Teflon) of the diluter system to eventually become saturated with HBCD. As this process progressed, an equilibrium was established. The result of this new equilibrium was that concentrations of HBCD in the dilution water were approximately the solubility of HBCD in well water under flow-through conditions.

Dissolved oxygen concentrations of > or = 97% of saturation were observed throughout the test. Water pH ranged from 8.1-8.4. Total organic carbon in the dilution water at test initiation was <1.0 mg C/L.

Daily observations during the test showed that daphnids in the negative control and solvent control groups appeared healthy and normal. With the exception of one aberrant mortality in the 4.6 ug/L (nominal) treatment group, all daphnids in all treatment groups appeared normal throughout the test with no mortalities or overt signs of toxicity. Based on these results, EC50 values for 24 and 48 hours were estimated to be > 6.8 ug/L (nominal), the highest concentration tested.

Conclusions

HBCD was not acutely toxic to *Daphnia magna*. The 48-hour EC50 value for daphids exposed to HBCD was > 6.8 ug/L (nominal) (>3.2 ug/L mean measured concentration), the highest concentration tested and twice HBCD's water solubility (3.4 ug/L). Based on the mortality, immobility and observation data, the 48-hour no mortality/immobility concentration and the no-observed-effect concentration was 6.8 ug/L (nominal) (3.2 ug/L mean measured concentration).

Data Quality

Reliability High

Data Reliability Remarks

This study was performed according to current guidelines and Good Laboratory Practices by a laboratory with considerable experience with these studies. Extensive attention was paid to analytical method development and performance.

Reference

>> Remarks

Graves, W and Swigert, J. (1997) Hexabromocyclododecane (HBCD): a 48-hour flow-through acute toxicity test with the cladoceran (*Daphnia magna*). Project Number: 439A-102. Wildlife International Ltd., Easton, MD.

General

EPA High Production Volume (HPV) Track

Ecotoxicity End Point:
Acute Toxicity to Aquatic Invertebrates

Sponsor ID	1103024	Albemarle Corporation	Create Date	1/6/01
CAS Number	3194156	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1103012	CMA Brominated Flame Retardant Industry Panel (BFRiP)	Completed:	N

Study sponsored by the Chemical Manufacturers Association Brominated Flame Retardant Industry Panel.

EPA High Production Volume (HPV) Track

Ecotoxicity End Point :
Toxicity to Aquatic Plants

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/9/01
CAS Number	3194550	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OECD Method 201; TSCA Title 40, CFR, Part 797, Section 1050

>> Test Type

static

>> GLP

Yes

>> Year study performed

1997

>> Species

Selenastrum capricornutum

>> End Point

Cell densities and area under the growth curve.

>> Analytical monitoring

HPLC/UV/VIS Detector; LOQ=0.400 ug/L

>> Exposure period

96 Hours

>> Statistical Method

Shapiro Wilk's; Bartlett's; Dunnett's; Bonferroni's t

Remarks for Method

The freshwater alga, *Selenastrum capricornutum*, was exposed to one of five test concentrations, a solvent control (DMF) or the negative (culture medium) control under static conditions for 96 hours. Three replicate test chambers were maintained for each treatment and control group. Nominal test concentrations were based on the solubility of the test substance in water (3.4 ug/L) and the results of an exploratory range finding toxicity test. Nominal test concentrations were 1.5, 2.2, 3.2, 4.6 and 6.8 ug HBCD/L. The highest dose tested was also confirmed analytically (HPLC with UV/VIS

EPA High Production Volume (HPV) Track

Ecotoxicity End Point :
Toxicity to Aquatic Plants

Sponsor ID	1100071	Albemarle Corporation	Create Date	4/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

detector).

Test solutions were inoculated with 1.0 mL of an inoculum with an approximate (~) density of 1.0×10^6 cells/mL to achieve a final cell density of 1.0×10^4 cells/mL. Samples of the test solutions were collected from each replicate test chamber at ~24 hour intervals during the test to determine cell density. Cell densities and area under the growth curve values were determined for each replicate and were used to calculate % inhibition values relative to the controls over the 96-hour exposure period. EC10, EC50 and EC90 values were calculated, if possible, based on cell densities and area under the growth curve values for each 24-hour interval. The no-observed-effect concentration (NOEC) was determined based on statistical evaluation of the cell densities and area under the growth curve values.

A primary stock solution was prepared by dissolving HBCD in dimethylformamide (DMF). The concentration of the stock was 0.068 mg HBCD/mL. Stock concentrations and the resultant test concentrations were prepared on a total product basis. A solvent control was prepared by diluting 250 μ L DMF to 2.5 L with culture medium to yield a solvent concentration equivalent of that in the treatment groups.

Original cultures of the freshwater algae, *Selenastrum capricornutum*, were obtained from UTEX - The Culture Collection of Algae at the University of Texas at Austin, and have been maintained in culture medium at Wildlife International Ltd, Easton, MD. Algal cells used in this test were obtained from Wildlife International Ltd cultures that had been actively growing in culture medium for a least two weeks prior to test initiation. The control organisms were expected to exhibit exponential growth over the 96-hour exposure period. Exponential growth phase, defined as the period of growth where the algal cells are dividing at a constant rate, is indicated by the linear section of the growth curve.

The algal cells were cultured and tested in freshwater algal medium. Test chambers were sterile 250-ml Erlenmeyer flasks plugged with foam stoppers, and containing 100 mL of test or control algal medium. The test chambers were shaken continuously at 100 rpm, and held in an environmental chamber at 24 ± 2 degrees C. Cool-white fluorescent lighting was used throughout the test (4310 \pm 431 lux). Samples of ~ 2 mL were collected from each treatment and control vessel at ~ 24 hour intervals during the 96-hour exposure. Cell counts were performed using an electronic particle counter (Coulter Electronics, Inc.). Samples of the test medium (test samples) were collected from each treatment and control group at the beginning and end of the test to measure concentrations of the test substance.

Results

>> Nominal concentration 0, 0.0015, 0.0022, 0.0032, 0.0046, 0.0068 mg/L

>> Measured concentration Negative control, Solvent control, and 0.0037 mg/L

>> Precision >

EPA High Production Volume (HPV) Track

Ecotoxicity End Point :
Toxicity to Aquatic Plants

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3194-55-5	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

>> Endpoint Type EC0

>> Endpoint Value 0

>> Unit used mg/L

>> Concentration Type Nominal

>> Endpoint Time 96

>> NOEC Precision >

>> NOEC 0

>> Unit used mg/L

>> NOEC Concentration Type Nominal

>> NOEC Effect(s) assesse cell density and growth

>> LOEC Precision >

>> LOEC 0

>> Unit used mg/L

>> LOEC Concentration Type Nominal

>> LOEC Effect(s) assesse cell density and growth

>> Response of Control Group (was it satisfactory?) Yes

>> Statistical results

No statistically significant differences ($p > 0.05$) were found between control and treated groups. Algal growth was not inhibited by HBCD, and EC10, EC50 or EC90 values could not be defined.

Results Remark

HBCD's 96 hr effect concentration in the freshwater algae tested could not be determined. The NOEC is greater than HBCD's water solubility.

The mean measured concentration at the 6.8 ug/L dose level was 3.7 ug/L.

Conclusions

HBCD's 96 hr effect concentration in the freshwater algae tested could not be determined. The NOEC is greater than HBCD's water solubility.

Data Quality

Reliability High

Data Reliability Remarks

EPA High Production Volume (HPV) Track

Ecotoxicity End Point :
Toxicity to Aquatic Plants

Sponsor ID	1103031	Albemarle Corporation	Create Date	4-8-01
CAS Number	3194558	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	110-012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

This study was performed according to current guidelines and Good Laboratory Practices by a laboratory with considerable experience with these studies. Extensive attention was paid to analytical method development and performance.

Reference

>> Remarks

C. Roberts and J. Swigert. Hexabromocyclododecane (HBCD) A 96-Hour Toxicity Test with the Freshwater Alga (*Selenastrum capricornutum*). Wildlife International Ltd. Project Number: 439A-103. June 3, 1997. Wildlife International Ltd., Easton, MD.

General

EPA High Production Volume (HPV) Track

Ecotoxicity End Point :
Toxicity to Aquatic Plants

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/8/01
CAS Number	3194556	Cyclododecano, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

Revision Date:

12/5/01

Test Substance

Remarks

A sample of the hexabromocyclododecane (HBCD) commercial product was obtained from one manufacturer, Great Lakes Chemicals Corporation (West Lafayette, IN).

Chemical Category

Method

>> Method/Guideline followed

Not specified.

>> Test Type

Not specified.

>> GLP Unknown

>> Year study performed 1987

>> Species

Skeletonema costatum, Thalassiosira pseudonana, Chlorella sp.

>> End Point cell numbers

>> Analytical monitoring Capillary column GLC; DL not specified.

>> Exposure period 72 Hr S. Costatum, T. Pseudonana; 96 Hr Chlorella

>> Statistical Method None - used linear regression to determine EC50

Remarks for Method

Each test was replicated. Population density was estimated by cell counts on a hemacytometer. The test article was introduced into growth flasks by adding 0.05 ml test article in acetone to 51 ml growth medium with algae. Algal species tested were S. costatum (Greville) Cleve, T. pseudonana Hasle and Heindal, and Chlorella sp., and were obtained from University of Rhode Island, Woods Hole Oceanographic Institution, and the Culture Collection of Algae, University of Texas at Austin, respectively. Growth media were prepared from seawater collected from an inshore site on the Gulf of Mexico and from five commercial sea salt formulations. Toxicity was expressed as the EC50 based on the cell numbers after incubation for 72 (S. costatum) or 96 hrs (T. pseudonana, Chlorella

EPA High Production Volume (HPV) Track

Ecotoxicity End Point :
Toxicity to Aquatic Plants

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3194550	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

sp.). The EC50 was derived by straight line graphical interpolation without calculation of confidence intervals. The highest test article concentration was determined by adding the test article slowly to growth media and observing the highest concentration at which crystals did not form.

Results

>> Nominal concentration

>> Measured concentration

>> Precision

>> Endpoint Type

>> Endpoint Value 0

>> Unit used

>> Concentration Type

>> Endpoint Time 0

>> NOEC Precision

>> NOEC 0

>> Unit used

>> NOEC Concentration Type

>> NOEC Effect(s) assesse

>> LOEC Precision

>> LOEC 0

>> Unit used

>> LOEC Concentration Type

>> LOEC Effect(s) assesse

>> Response of Control Group (was it satisfactory?)

>> Statistical results

Results Remark

EPA High Production Volume (HPV) Track

Ecotoxicity End Point :
Toxicity to Aquatic Plants

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/3/01
CAS Number	3124556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	N

Growth of *Chlorella* sp. was not inhibited by HBCD at the highest dose tested, 1.5 mg/L. HBCD's EC50 for *S. costatum* ranged from 9.0-12.0 ug/L in the six media. Similarly, HBCD's EC50 in *T. pseudonana* ranged from 0.05-0.37 mg/L in the six media.

The pH of the six different growth media ranged from 7.6-8.2. No relationship of pH to toxicity was found for HBCD.

There was little variation in the response of *S. costatum* to HBCD among the media, but the response to *T. pseudonana* varied widely. *S. costatum* may be more sensitive to HBCD than *T. pseudonana*.

Conclusions

HBCD's 96 hour EC50 in *Chlorella* sp., tested in 6 different growth media, was > 1.5 mg/L. HBCD's 72 hour EC50 in *S. Costatum* and *T. Pseudonana* in 6 different growth media ranged from 0.009-0.012 and 0.5-0.36 mg/L, respectively. All EC50 values determined in the three marine algae were greater than HBCD's water solubility (0.0034 mg/L).

Data Quality

Reliability good

Data Reliability Remarks

Reference

>> Remarks

Walsh, G., Yoder, M., McLaughlin, L., Lores, E. (1987) Responses of marine unicellular algae to brominated organic compounds in six growth media. *Ecotoxicology and Environmental Safety*, 14, 215-222.

General

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	Y

Revision Date:

4/10/01

Test Substance

Remarks

A form of hexabromocyclododecane (HBCD) supplied as test article by Saytech Inc. No further details are available.

Chemical Category

Method

>> Method/Guideline followed

Not known.

>> GLP Unknown

>> Year study performed 1978

>> Species

rat

>> Strain no data

>> Sex Both

>> Number of males per dose

5

>> Number of females per dose

5

>> Vehicle Corn oil

>> Route of Administration

Oral

Remarks for Method

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	11/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Albino rats in groups of ten (5M:5F), 192-260 g, were administered a single dose (10 g/kg) orally and observed for 14 days. The highest volume used was 40 ml/kg. The vehicle was corn oil

Results

>> Precision

>

>> Acute Lethal Value

10000

>> Unit

mg/kg-bw

>> Deaths per Dose

One of five males died on test. No females died on test.

Results Remark

Conclusions

The oral LD50 of HBCD in the rat was > 10,000 mg/kg body weight.

Data Quality

Reliability

Acceptable

Data Reliability Remarks

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	3/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

This study is old and not performed according to current guidelines. Nonetheless, the results are consistent with the general lack of toxicity associated with this material in other mammalian studies. Thus, it was found acceptable.

Reference

>> Remarks

Lewis, C and Palanker, A. (1978) Final Report. Oral LD50 (Rat). Experiment Reference No.: 78385-1. Consumer Product Testing Company Incorporated, Fairfield, NJ.

General

Sponsored by Saytech, Inc., Sayreville, NJ.

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	1191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1301012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

12/5/01

Test Substance

Remarks

A form of hexabromocyclododecane (HBCD) supplied as test article by Saytech Inc. No further details are available.

Chemical Category

Method

>> Method/Guideline followed

Not known.

>> GLP

Unknown

>> Year study performed

1978

>> Species

rabbit

>> Strain

New Zealand White

>> Sex

Both

>> Number of males per dose

3

>> Number of females per dose

3

>> Vehicle

None

>> Route of Administration

Dermal

Remarks for Method

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	Y

Method was described as that of Hagen (1959). Albino rabbits in groups of six (3M:3F), one half with abraded skin, 1.88-2.07 kg, highest dose level mechanically possible, single application dermally under occluded patch, observed for 14 days. Material used as received. Upper limit possible due to mechanical and physical limitations is 8 g/kg body weight.

Results

>> Precision

>

>> Acute Lethal Value

8000

>> Unit

mg/kg-bw

>> Deaths per Dose

No animals died on test.

Results Remark

Conclusions

The dermal LD50 of HBCD in rabbits was > 8,000 mg/kg body weight.

Data Quality

Reliability

Acceptable.

Data Reliability Remarks

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3101556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

This study is old and not performed according to current guidelines. Nonetheless, the results are consistent with the general lack of toxicity associated with this material in other mammalian studies. Thus, it was found acceptable.

Reference

>> Remarks

Lewis, C and Palanker, A. (1978) Final Report. Dermal LD50 (Rabbit). Experiment Reference No.: 78385-2. Consumer Product Testing Company Incorporated, Fairfield, NJ.

General

Sponsored by Saytech, Inc., Sayreville, NJ.

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	1191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	3
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

4/11/01

Test Substance

Remarks

A form of hexabromocyclododecane (HBCD) supplied as test article by Saytech Inc. No further details are available.

Chemical Category

Method

>> Method/Guideline followed

Not known.

>> GLP

Unknown

>> Year study performed

1978

>> Species

rat

>> Strain

no data

>> Sex

Both

>> Number of males per dose

5

>> Number of females per dose

5

>> Vehicle

None

>> Route of Administration

Inhalation

Remarks for Method

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	1100621	Albemarle Corporation	Create Date	1/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Albino rats in groups of 10 (5M:5F), 233-292 g, exposed to concentrations of 200 mg/L (highest possible chamber concentration) for one hour, observed two weeks. Material used as received.

Results

>> Precision >

>> Acute Lethal Value 200

>> Unit mg/L(air)

>> Deaths per Dose

No animals died on test.

Results Remark

Conclusions

The inhalation LC50 of HBCD in rats was > 200 mg/L for a 1 hour exposure.

Data Quality

Reliability Acceptable.

Data Reliability Remarks

EPA High Production Volume (HPV) Track

Toxicity End Point:
Acute Toxicity

Sponsor ID	100021	Albemarle Corporation	Create Date	11-01
CAS Number	319156	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	3
Consortia ID	101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

This study is old and not performed according to current guidelines. Nonetheless, the results are consistent with the general lack of toxicity associated with this material in other mammalian studies. Thus, it was found acceptable.

Reference

>> Remarks

Lewis, C and Palanker, A. (1978) Final Report. Inhalation LC50 (Rat). Experiment Reference No.: 78385-2. Consumer Product Testing Company Incorporated, Fairfield, NJ.

General

Sponsored by Saytech, Inc., Sayreville, NJ.

EPA High Production Volume (HPV) Track

Toxicity End Point:
Developmental Toxicity/Teratogenicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	11/6/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	<input type="checkbox"/>

Revision Date:

12/5/01

Test Substance

Remarks The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method >> Method/Guideline followed

EPA OPPTS Method 870.3700; OECD 414

>> GLP Yes

>> Year study performed 1999

>> Species

rat

>> Strain Mammal strain Sprague-Dawley

>> Sex F

>> Number of males per dose 0

>> Number of females per dose 25

>> Route of Administration Oral

>> Days of Gestation 6-19

>> Frequency of treatment Once daily

>> Doses 0, 250, 500, 1000 mg/kg body weight

>> Control Group Yes

Concurrent control

>> Statistical Method

See Remarks for Method.

Remarks for Method

EPA High Production Volume (HPV) Track

Toxicity End Point:
Developmental Toxicity/Teratogenicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	

Hexabromocyclododecane (HBCD) was administered by gavage in corn oil to three groups of 25 bred Crl:CD(SD)IGS BR (Charles River Laboratories, Raleigh, NC) rats once daily from gestation days 6 through 19. Dosage levels were 250, 500 and 1000 mg/kg/day administered in a dose volume of 5 ml/kg. A concurrent control group composed of 25 bred females received the vehicle, corn oil, on a comparable regimen. Clinical observations, body weights and food consumption were recorded. On gestation day 20, a laparohysterectomy was performed on all animals. The uteri and ovaries were examined and the numbers of fetuses, early and late resorptions, total implantations and corpora lutea were recorded. Mean gravid uterine weights and net body weight changes were calculated for each group. The fetuses were weighed, sexed and examined for external soft tissue and skeletal malformations and variations.

Appropriate statistical tests were used for each end point and included a one-way ANOVA with Dunnett's test, and Kruskal Wallis test with Mann-Whitney U test.

Results

>> Maternal Precision/NOAEL	=	
>> Maternal NOAEL dose	1000	>> Unit used mg/kg-bw
>> Maternal NOAEL effect	None	
>> Maternal Precision/LOAEL	>	
>> Maternal LOAEL dose	1000	>> Unit used mg/kg-bw
>> Maternal LOAEL effect	None	
>> Developmental Precision/NOAEL	=	
>> Developmental NOAEL dose	1000	>> Unit used mg/kg-bw
>> Developmental NOAEL effect	None	
>> Developmental Precision/NOAEL	>	
>> Developmental LOAEL dose	1000	>> Unit used mg/kg-bw
>> Developmental LOAEL effect	None	
>> Actual dose	As given above.	
>> Maternal data with dose level (with NOAEL value).		

No adverse effects detected.

EPA High Production Volume (HPV) Track

Toxicity End Point:
Developmental Toxicity/Teratogenicity

Sponsor ID	1100621	Albemarle Corporation	Create Date	4/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	

>> Fetal data with dose level (with NOAEL value).

No adverse detected.

>> Statistical results

See Methods.

Results Remark

All maternal animals survived to the scheduled necropsy on gestation day 20. One female in the 500 mg/kg/day group delivered on gestation day 20 and was examined at the scheduled laparohysterectomy. No treatment-related clinical signs were observed at any dose level. Body weight gain and food consumption were not adversely affected at any dose level. At necropsy, no treatment-related findings were observed. Intrauterine growth and survival were unaffected by test article administration at any dose level. No treatment-related fetal malformations or developmental variations were observed in any of the treated groups.

Conclusions

The no-observed-adverse-effect level for maternal toxicity and developmental toxicity was 1000 mg HBCD/kg/day administered on days 6-19 of gestation.

Data Quality

Reliability

High

Data Reliability Remarks

This study was performed according to current guidelines for repeated dose studies under Good Laboratory Practices by a laboratory experienced in the performance of studies of this type.

Reference

>> Remarks

Stump, D. (1999) A Prenatal Developmental Toxicity Study of Hexabromocyclododecane (HBCD) in Rats. Laboratory Study No.: WIL-186009. WIL Research Laboratories, Inc., Ashland, OH.

General

Sponsored by Chemical Manufacturers Association Brominated Flame Retardant Industry Panel.

EPA High Production Volume (HPV) Track

Toxicity End Point:
Developmental Toxicity/Teratogenicity

Sponsor ID	1102021	Albemarle Corporation	Create Date	1/7/01
CAS Number	1194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	

Revision Date:

12/5/01

Test Substance

Remarks The test article was manufactured by Daiichi Kogyo Seiyaku K.K. No further information on its composition is known.

Chemical Category

Method >> Method/Guideline followed

Not specified.

>> GLP Unknown

>> Year study performed 1985

>> Species

rat

>> Strain Mammal strai Wistar

>> Sex F

>> Number of males per dose 0

>> Number of females per dose 20

>> Route of Administration Oral

>> Days of Gestation 0-20

>> Frequency of treatment Daily

>> Doses 0, 0.01, 0.1 and 1% of the Diet

>> Control Group Yes Concurrent control

>> Statistical Method

Not specified.

Remarks for Method

EPA High Production Volume (HPV) Track

Toxicity End Point:
Developmental Toxicity/Teratogenicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	<input type="checkbox"/>

The Murali et al study consisted of a 7 day dose range finding study (n=5 rats/dose group) and a combined teratogenicity-developmental study (n=20/dose group). Doses in the 7 day range finding study were 0, 0.3, 1, 3 or 10 g/kg/day. Doses as high as 10 g/kg/day produced no evidence of toxicity. A statistically significant ($P < 0.01$) increase in liver weight was noted in groups receiving > 1 g/kg/day. Doses for the combined teratogenicity-developmental study were based on this increase in liver weight.

In the combined teratogenicity-developmental study, pregnant female rats were fed diets containing 0, 0.01, 0.1, or 1% HBCD on days 0-20 of gestation. Daily doses were estimated by the authors to be 0, 5, 50 or 500 mg/kg/day and the average total dose/rat/group was estimated to be 0, 0.13, 1.28 or 12.0 g/kg. Rats were observed daily and body weight and food consumption measured. Fourteen rats from each group were sacrificed on day 20 of gestation and their fetuses were examined for toxicity or teratogenicity. Approximately 150 fetuses/dose level were examined for evidence of teratogenicity. All fetuses from all litters were examined for signs of external anomalies. Approximately 2/3 of the fetuses/dam were examined for skeletal abnormalities; the remaining fetuses from each dam were examined for any abnormalities of the internal organs. In addition, six rats from each group were allowed to deliver their litters and growth of the litters was observed until the 7th week post-parturition.

Results

>> Maternal Precision/NOAEL	>	
>> Maternal NOAEL dose	1000	>> Unit used mg/kg in feed
>> Maternal NOAEL effect	No adverse effects, increased liver wt at 1% dose.	
>> Maternal Precision/LOAEL	>	
>> Maternal LOAEL dose	1000	>> Unit used mg/kg in feed
>> Maternal LOAEL effect	No adverse effects.	
>> Developmental Precision/NOAEL	>	
>> Developmental NOAEL dose	1000	>> Unit used mg/kg in feed
>> Developmental NOAEL effect	No adverse effects.	
>> Developmental Precision/NOAEL	>	
>> Developmental LOAEL dose	1000	>> Unit used mg/kg in feed
>> Developmental LOAEL effect	No adverse effects.	
>> Actual dose	Estimated as 0, 5, 50, 500 mg HBCD /kg bd wt/day	
>> Maternal data with dose level (with NOAEL value).		

EPA High Production Volume (HPV) Track

Toxicity End Point:
Developmental Toxicity/Teratogenicity

Sponsor ID	1100021	Aibemarle Corporation	Create Date	1/6/01
CAS Number	1193156	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101812	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	

Only effect detected in dams was an increase in liver weight at the 1% dose level.

>> Fetal data with dose level (with NOEL value).

No effects detected in fetuses.

>> Statistical results

See results remarks.

Results Remark

The authors' estimated the doses in the feed were equivalent to 0, 5, 50 or 500 mg HBCD /kg body weight /day. No adverse effects were detected in any treatment group with respect to maternal weight gain, food consumption, or gross appearance of internal organs. The mean liver (absolute and relative to body weight) weight in the 1% group was statistically different (higher) from the control mean. Normal development was seen in neonates carried through to six weeks of age.

There was no adverse effect of treatment on the number of corpora lutea, implants, resorptions, live fetuses, sex ratio, or body or placental weight. No fetal deaths occurred in any group. No external, skeletal or visceral malformations were detected. A few skeletal variations were detected but where of similar types and numbers in the control and treated groups.

There was no significant differences between the control and treated groups in the number of implantation, live newborns, dead newborns, live newborn parturition index. The weaning and survival index was comparable in the control and treated groups. Body weight changes in the newborns was comparable in all groups.

Conclusions

No reproductive or developmental effects were detected in rats at HBCD doses up to 1% in the diet (~500 mg/kg/d) administered from days 0-20 of gestation. Further, normal development was seen in neonates carried through to six weeks of age.

Dose levels: 0, 0.01, 0.1, or 1% HBCD on days 0-20 of gestation [Murai estimate: 0, 5, 50 or 500 mg/kg/day]. No teratogenic effects. Normal development in neonates carried through age 6 wks. NOEL = 1% of diet.

Data Quality

Reliability

Good.

Data Reliability Remarks

EPA High Production Volume (HPV) Track

Toxicity End Point:
Developmental Toxicity/Teratogenicity

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	

One author of this study was associated with the National Institute of Hygienic Science, Osaka branch.

Reference

>> Remarks

Murai, T. Kawasaki, H., Kanoh, S. 1985. Studies on the toxicity of insecticides and food additives in pregnant rats - Fetal toxicity of Hexabromocyclododecane. Pharmacometrics (Japan) 29(6):981-986.

General

Funding for this study was provided by Japan's Ministry of Health and Welfare.

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	31915-99	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OECD Method 407

>> GLP Yes

>> Year study performed 1997

Were Good Laboratory Practices followed in the st

>> Species

rat

>> Strain Mammal strai Sprague-Dawley

>> Sex Both

>> Number of males per dose

6

>> Number of females per dose

6

>> Route of Administration

>> Exposure Period

Duration of study in days (for example, 28 days, 90 d)

>> Frequency of treatment

Number of doses per day, week, etc. This is particularly relevant for inhalation experiments -- 6hrs/day, 5 d

>>Doses

List all doses used in te

>> Control Group

Concurrent contro

>> Post observation period

Length of time animals observed after last d

>> Statistical Method

Cite statistical methods use

Remarks for Method

Hexabromocyclododecane (HBCD) was administered orally by gavage in corn oil to three groups of Sprague-Dawley Cri:CD BR (Charles River Laboratories, Inc., Portage, MI) rats for a period of 28 consecutive days at doses of 125, 350 or 1000 mg/kg/day administered in a dosage volume of 5 ml/kg. The test groups consisted of 6 males and 6 females in the 125 and 350 mg/kg/day groups, and 12 males and 12 females in the 1000 mg/kg/day group. A concurrent control group (n=12 males and females) was treated in a similar manner with the vehicle, corn oil. At the end of the dosing period, 6 animals/sex/group were euthanized and necropsied. The remaining 6 animals/sex in the control and 1000 mg/kg/day groups remained on test untreated for a 14 day recovery period. At the end of the recovery peroid, all animals were euthanized and necropsied. Animals were 6 weeks of age at study initiation.

Animals were observed twice daily for mortality and morbundity. Clinical signs were recorded daily. Body weights and food consumption were measured weekly. Functional observational battery and motor activity evaluations were performed during weeks 1 (pretest), 3, and 5 (recovery). Samples for hematology and serum chemistry evaluations were collected at the primary (28 day) and recovery (42 day) necropsies. Complete necropsies were performed on all rats. The brain, liver, kidney, heart, spleen, testes and epidymus or ovaries, adrenal glands, and thymus from all animals were weighted at each necropsy. Approximately 40 tissues were

Results

collected and preserved at each necropsy from each animal. The following tissues were examined microscopically from the control and high dose animals: liver, kidney, heart, spleen, testes (males), prostate (males), seminal vesicles (males), epididymus (males), ovaries (females), adrenal glands, thymus, bone with marrow (sternebra), brain, stomach, cecum, duodenum, ileum, jejunum, lymph node, peripheral nerve (sciatic), spinal cord, lung, trachea, uterus (females), urinary bladder, and all gross lesions. The lungs, liver, kidney, stomach, gross lesions and target organs were examined in all dose levels.

Body weights, weight gain, food consumption, functional observation battery and motor activity results of treated animals were compared statistically by sex and treatment day to their respective control groups ($p < 0.05$ or < 0.01).

Concentrations of the dosing suspensions were confirmed. Homogeneity determinations were performed on study days 0, 13, and 27.

All statistical analyses were conducted using two-tailed tests for minimum significance levels of 1% and 5% comparing the treatment groups to the vehicle control group by sex. Analysis of body weight change, food consumption, clinical pathology values, continuous functional observational battery data and absolute and relative organ weight data were analyzed with a one-way analysis of variance followed by Dunnett's test. Discontinuous (ordinal or descriptive) functional observational battery data were analyzed using Fisher's exact test. Statistical tests for locomotor activity data were performed using SAS/STAT statistical software. Clinical laboratory values for cell types that occur at a low incidence (i.e., monocytes, eosinophils and basophils) were not subjected to statistical analysis.

>> NOAEL Precision

>=

>> NOAEL dose

1000

>> Unit

mg/kg-bw

>> NOAEL Effec

Increase in liver weight in the absence of histopathologic or clinical chemistry changes.

(e.g., decrease in body weight, organ

>> LOAEL Precision

>

>> LOAEL dose

1000

>> Unit

mg/kg-bw

>> LOAEL Effect

(e.g., decrease in body weight, organ

None noted.

>> Actual dose received by dose level by sex (if available)

Test article administered by gavage.

>> Toxic response A brief narrative describing toxic response or effects, by dose

No evidence of toxicity was observed at any dose level.

>> Statistical results Note statistical results, with appropriate ρ value

See Results Remarks section.

Results Remark Provide at a minimum qualitative descriptions of elements where dose effect related observations of elements were seen:

Survival was not affected by administration of the test article. All animals survived to the scheduled necropsy. Clinical signs observed during the study were nonspecific, low in incidence, non-dose-related, and not considered related to test article.

Body weights, weight gain and food consumption were not affected by treatment. No statistically significant differences in mean body weight between control and treated animals were detected with the exception of an increase in mean female body weight in the 350 mg/kg/day group during week 2. Mean female body weight at that time point was 196 g in the 350 mg/kg/day group vs. 179 g in the control group. No statistically significant differences in body weight gain between the control and treated animals with the expectation of a decrease in mean male body weight gain in the 1000 mg/kg/day recovery group during week 1 of recovery. Mean male body weight gain at that time point was 21 g vs 31 g in the control group; mean male body weight was not statistically different from the control mean. No statistically significant differences in food consumption between control and treated animals were detected with the exception of an increase in mean female food consumption in the 350 mg/kg/day during weeks, -1, 1, and 2 of treatment. Mean female food consumption at those time points were 18, 17 and 17 g vs. 16, 15 and 15 g in the control group, respectively.

Results of the functional observation battery and motor activity tests were not affected by treatment. No statistically significant differences were observed between the control and

treated animals at any time point ($p < 0.05$).

No statistically significant differences between control and treated animals were found for hematology parameters with the exception of an increase in mean activated partial thromboplastin time in the 1000 mg/kg/day males on week 4 and a decrease in the mean prothrombin time in the 1000 mg/kg/day females on week 4. These statistical differences were not of toxicological significance.

No toxicologically significant effects on serum chemistry values related to test article administration were observed at the 28 day primary and 42 day recovery necropsies. Scattered instances of statistically significant differences between treated and control animals were detected for some serum chemistry parameters at the 28 day primary necropsy. These scattered statistical differences were not considered toxicologically significant because the statistical differences occurred in the absence of a dose response, in the absence of the accompanying clinical chemistry changes expected, in the opposite direction from what occurs in a toxic state, in a direction which is without physiologic significance, or due to potential interference with the laboratory method. No statistically significant differences in serum chemistry parameters were detected between groups at the 42 day recovery necropsy.

No gross lesions attributable to test article administration were detected at either necropsy. Gross lesions were nonspecific, low in incidence, non-dose-related, and considered incidental.

No microscopic lesions attributable to test article administration were detected on histopathologic exam. Microscopic changes were nonspecific, low in incidence, non-dose-related and considered incidental.

No statistically significant differences in organ weight or organ to body weight ratios were detected between control and treated animals with one exception. Absolute liver weights were statistically significantly increased with respect to control mean at the 28 day necropsy in males in the 1000 mg/kg/day group and in females in the 350 and 1000 mg/kg/day groups. Liver to body weight ratios in the 350 and 1000 mg/kg/day male and female groups were statistically increased at the 28 day necropsy. At the recovery necropsy, male absolute and liver to body weight ratio were statistically comparable to the control mean. Female absolute liver weight and liver to body weight ratio were statistically increased compared to the control mean. The difference in absolute liver weight between control and treated females was less pronounced at the end of the recovery period, indicating the increase in liver weight was reversible in females as well as males. In the absence of test article related histologic and serum chemistry changes, increases in liver weight are considered an adaptive rather than toxic response, are not uncommon in the rat, and are most likely the result of microsomal

induction.

Conclusions

Input optional information or further explain the contents of a particular section, much as is done in the "Discussion" portion of a publication in academic journals.

No systemic toxicity was observed at any dose level. The No Observed Adverse Effect Level of HBCD administered orally to male and female rats for 28 consecutive days was ≥ 1000 mg/kg/day, the highest dose tested.

Data Quality

Reliability High

Denote the reliability of data, at the discretion of the person preparing the robust su

Data Reliability Remarks Add comments about how reliability of data was determined, or add re

This study was performed according to current guidelines for repeated dose studies under Good Laboratory Practices by a laboratory experienced in the performance of studies of this type.

Reference

Cite the full reference for the critical study on which the robust study summary is based. List other appropriate references that support this summary and, have been reviewed for

>> Remarks

Chengelis, C. (1997) A 28-Day Repeated Dose Oral Toxicity Study of HBCD in Rats. Laboratory Study Number: WIL-186004. WIL Research Laboratories, Inc., Ashland, OH.

General

Add any statement that doesn't fit into any of the other f

Study sponsored by the Chemical Manufacturers Association Brominated Flame Retardant Industry Panel.

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3191-55-5	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1201012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	1/2



Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OECDMethod 408, EPA OPPTS Method 870.3100

>> GLP Yes

>> Year study performed 2001

Were Good Laboratory Practices followed in the st

>> Species

rat

>> Strain Mammal strain Sprague-Dawley

>> Sex Both

>> Number of males per dose 15

>> Number of females per dose 15

>> Route of Administration Oral

>> Exposure Period

Duration of study in days (for example, 28 days, 90 d)

>> Frequency of treatment

Number of doses per day, week, etc. This is particularly relevant for inhalation experiments – 6hrs/day, 5 d

>>Doses

List all doses used in te

>> Control Group

Concurrent contro

>> Post observation period

Length of time animals observed after last d

>> Statistical Method

Cite statistical methods use

Remarks for Method

The test article, a composite of three lots of commercial hexabromocyclododecane (HBCD), was administered by oral gavage in corn oil once daily to four groups of Crl:CD(SD)IGS BR rats (n=15/sex/group) at dose levels of 0 (control), 100 (low), 300 (mid) and 1000 (high) mg/kg/day seven days per week for 90 days. The dosage volume was 5 ml/kg. The control animals received the vehicle, corn oil, only. At the end of the 90-day treatment period, 10 animals/sex/group were euthanized and necropsied. The remaining rats continued on test untreated for a 28-day recovery period prior to necropsy.

Results

In addition to the main toxicology groups, two satellite groups of 20 animals/sex/group were treated concurrently in an identical manner at dose levels of 0 or 1000 mg HBCD/kg/day for up to 90 days. Body weights were recorded weekly. Two animals/sex/group were euthanized on study days 2, 6, 9, 13, 20, 27, 55, 89, 104 and 118, and blood and body fat (mesenteric and/or omental) were collected. The body fat was analyzed for HBCD content.

Animals in the main toxicology groups were observed twice daily throughout the study for mortality and morbidity. Body weights and food consumption were measured weekly. Blood was collected at study weeks 3 (n=5/sex/group), 13 (n=10/sex/group) and 17 (n=5/sex/group) for hematology, serum chemistry and hormone (T3, T4 and TSH) measurements. Urine was collected prior to each necropsy, at study weeks 13 and 17, for urinalysis. Ocular examinations were performed prior to study initiation and during study weeks 12 and 15.

Functional Observational Battery and Locomotor Activity evaluations were performed on 5 animals/sex/group prior to study initiation, during the last week of test article administration (study week 13), and during the recovery period. An examination of vaginal cytology (for estrus cycle determinations) was performed on study days 69-90. At each necropsy, sperm motility/viability, morphology, and number were assessed. Complete necropsies were performed on all animals. Approximately 40 organs or tissues/animal were collected and preserved. The adrenals, brain, epididymides, heart, kidneys, liver, ovaries, prostate, spleen, testes, thymus, thyroids with parathyroids, and uterus with cervix were weighed. Paraffin sections of tissues stained with hematoxylin and eosin from the control and 1000 mg/kg/day dose groups and the liver, lungs and thyroid glands in the 100 and 300 mg/kg/day doses, and gross lesions from all animals were examined under the light microscope. Livers from five randomly chosen animals/sex from the control and 1000 mg/kg/day dose groups were examined microscopically using Oil Red O or periodic acid Schiff's (PAS) reagent for evidence of lipid accumulation or glycogen accumulation/depletion, respectively. Statistical comparisons by sex and treatment day were made between the control and treated animals where indicated ($p < 0.05$).

>> NOAEL Precision

>=

>> NOAEL dose

1000

>> Unit

mg/kg-bw

>> NOAEL Effect

See Results Remarks.

(e.g., decrease in body weight, organ

>> LOAEL Precision

>

>> LOAEL dose

1000

>> Unit

mg/kg-bw

>> LOAEL Effect

No adverse effects detected.

(e.g., decrease in body weight, organ

>> Actual dose received by dose level by sex (if available)

As given under Doses.

>> Toxic response A brief narrative describing toxic response or effects, by dose

See Results Remarks.

>> **Statistical results** Note statistical results, with appropri ρ value

See Results Remarks.

Results Remark Provide at a minimum qualitative descriptions of elements where dose effect related observations of elements were seen:

No test article-related effect on mortality occurred. Clinical signs were non-specific, low in incidence, non-dose-related and not related to test article administration. No test article-related changes occurred in body weight, food consumption, Functional Observational Battery or Locomotor Activity. No test article-related effects on hematologic parameters were noted. No test article-related ocular lesions were detected at the ophthalmic exams. No test article-related changes were noted on the estrus cycle as determined by vaginal cytology, or on sperm motility/viability, morphology, and number. Instances of statistically significant differences between control and some treatment groups were detected at study week 13 in the clinical chemistry data, hormone data, organ weight data and histology findings. They were generally secondary to the inducing effects on the liver or were otherwise not considered adverse effects of treatment as discussed further below.

Statistically significant ($p < 0.05$) test article-related clinical chemistry changes at week 13 include an increase in albumin (all dose levels for males), total protein (all dose levels for females and 1000 mg/kg/day for males), globulin (300 and 1000 mg/kg/day for females), and chloride (all doses for both sexes). In addition, increased gamma glutamyltransferase levels were noted in the 1000 mg/kg/day group ($p < 0.05$). Thyroxine (T4) levels were decreased at study week 13 compared to the control mean in all male dose groups and the 300 and 1000 mg/kg/day dose females ($p < 0.05$). There were no corresponding statistical effects on T3 and TSH. While potentially test article-related, the changes in serum chemistry parameters were not of sufficient magnitude to be adverse, occurred in otherwise clinically normal animals, tended to be within or close to historical control values, and were not present at the end of the recovery period; furthermore, these serum albumin and gamma glutamyltransferase increases were probably secondary to the increases in liver weight. The increases in serum chloride were probably secondary to the presence of free bromide in the test article preparation which interfered with the chloride determination methodology. The decrease in T4, which was also reversible, was also probably secondary to increased liver weight (secondary to microsomal enzyme induction, known to cause increased metabolism and clearance of T4 in the rat).

The incidence of observations noted at gross necropsy was low and there was no evidence of frank organ damage. On histopathologic examination of tissues, relatively mild findings occurred in both the control and treated groups. Potential test article-related histologic changes were identified in the liver and thyroid glands but these would not be considered indicative of frank toxicity. These organs were examined microscopically in all groups at both necropsies. The liver changes in male rats at the 90-day necropsy (Study Week 13) were characterized as minimal hepatocellular vacuolation and occurred in 10% of control males and ~50% of the males at 100, 300 and 1000 mg/kg/day. Minimal hepatocellular vacuolation was also detected in females in the control and test article treated groups without a clear dose response (3 to 4/10 animals per group) but, mild and moderate vacuolation was detected in females only in the 300 (1/10) and 1000 mg/kg/day (2/10) dose groups. Minimal to mild hepatocellular hypertrophy was also detected only in the 1000 mg/kg/day group (5/10) females. Minimal thyroid follicular cell hypertrophy was detected 1/10, 1/10, 5/10 and 7/10 males in the control, 100, 300 and 1000 mg/kg/day groups, respectively and in 4/10 and 3/10 females in the 300 and 1000 mg/kg/day groups respectively. In addition, mild thyroid follicular hypertrophy was detected in 4/10 females in the 1000 mg/kg/day group. The histologic changes in the liver were accompanied by an increase in liver weight. In contrast there were no statistically significant changes in thyroid weight (absolute, relative to body weight and relative to brain weight). At study week 13, mean liver weights in all dose levels of both sexes (absolute, relative to body weight and relative to brain weight) were increased compared to the male and female control means ($p < 0.05$). The increases in liver weight were a result of a microsomal enzyme inducing effect¹ and were not typically considered indicative of toxicity in absence of frank organ damage. The reversible histologic changes (vacuolation and hypertrophy) are often found to accompany increased liver weight caused by liver enzyme induction. At week 17, the liver changes (weight and histology) had at least partially, if not fully, resolved in all treated groups without delayed or long-term toxic effects. The histologic changes in the thyroid had also nearly completely resolved except in the 1000 mg/kg/day group females, where partial recovery occurred.

Increases in mean prostate weight were noted in the 1000 mg/kg/day group males at the primary necropsy. However, the increases in prostate weight were probably not of toxicological significance since the increases did not persist to the recovery period, there were no correlating histologic findings and no change in sperm production.

HBCD was detected in the adipose tissue of male and female rats treated with 1000 mg/kg/day for up to 90 days. Isomer-specific analysis showed that the relative isomer concentrations in adipose tissue at all time points were $\alpha > \gamma > \beta$ which is in contrast to the test article composition ($\gamma > \alpha > \beta$). Steady state levels were achieved by study day 27. Levels in male and female rats were similar at all time points and declined during the recovery period.

All the test article-related changes at 100 and 300 mg/kg/day were mild, reversible, generally secondary to hepatic enzyme induction (which is an adaptive not a toxic change) and without effect on the clinical condition of the animals. The additional findings observed at 1000 mg/kg/day (increased gamma glutamyltransferase and additional increases in the size of the liver and prostate), were also reversible, not associated with specific target organ damage or diminished function and were, therefore, probably of limited, if any, toxicologic significance. On this basis the no-observed-adverse-effect level (NOAEL) of HBCD administered to CrI:CD®(SD)IGS BR rats by gavage in corn oil for 90 days is 1000 mg/kg/day.

Conclusions

Input optional information or further explain the contents of a particular section, much as is done in the "Discussion" portion of a publication in academic journals.

The no-observed-adverse-effect level (NOAEL) of HBCD administered to CrI:CD®(SD)IGS BR rats by gavage in corn oil for 90 days is 1000 mg/kg/day, the highest dose tested.

Data Quality

Reliability High

Denote the reliability of data, at the discretion of the person preparing the robust su

Data Reliability Remarks Add comments about how reliability of data was determined, or add re

This study was performed according to current guideline under good laboratory practices by laboratory with considerable experience in this area.

Reference

Cite the full reference for the critical study on which the robust study summary is based. List other appropriate references that support this summary and, have been reviewed for

>> Remarks

Chengelis, C. An Oral (Gavage) 90 Day Toxicity Study of HBCD in Rats. Laboratory Study No. WIL-186012. WIL Research Laboratories, Inc., Ashland, Ohio. 2001.

General

Add any statement that doesn't fit into any of the other f

Sponsored by the American Chemistry Council's Brominated Flame Retardant Industry Panel (BFRIP).

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	319-850	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	3
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	YES

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a commercial HBCD product ("Hexabromid S") produced at one time by BASF in Germany. BASF no longer manufactures HBCD.

Chemical Category

Method

>> Method/Guideline followed

Not specified.

>> GLP

>> Year study performed

Were Good Laboratory Practices followed in the st

>> Species

>> Strain Mammal strai

>> Sex

>> Number of males per dose >> Number of females per dose

>> Route of Administration

>> Exposure Period

Duration of study in days (for example, 28 days, 90 d)

>> Frequency of treatment

Number of doses per day, week, etc. This is particularly relevant for inhalation experiments -- 6hrs/day, 5 d

>> Doses

List all doses used in te

>> Control Group

Concurrent contro

>> Post observation period

Length of time animals observed after last d

>> Statistical Method

Cite statistical methods use

Remarks for Method

HBCD ("Hexabromid S") was tested in Sprague-Dawley rats (10/sex/group) at doses of 0, 1, 2.5 and 5% of the diet for 28 days. Doses calculated from the actual body weights and food consumption in this study are 0, 940, 2410, and 4820 mg/kg body weight/day.

Results

>> NOAEL Precision

>> NOAEL dose >> Unit

>> NOAEL Effec

(e.g., decrease in body weight, organ

>> LOAEL Precision =

>> LOAEL dose 5000

>> Unit mg/kg in feed

>> LOAEL Effect
(e.g., decrease in body weight, organ

Decrease in body weight at a dose level of 5% in the diet.

>> Actual dose received by dose level by sex (if available)

0, 940, 2410, and 4820 mg/kg body weight/day

>> Toxic response A brief narrative describing toxic response or effects, by dose

See Results Remarks.

>> Statistical results Note statistical results, with appropriate p value

See Results Remarks.

Results Remark Provide at a minimum qualitative descriptions of elements where dose effect related observations of elements were seen:

HBCTD ("Hexabromid S") was tested in Sprague-Dawley rats (10/sex/group) at doses of 0, 1, 2.5 and 5% of the diet for 28 days. Doses calculated from the body weights and food consumption are 0, 940, 2410, and 4820 mg/kg body weight/day.

No clinical signs were observed at the 1% dose levels. No significant change in mean body weight between the control and the 1 and 2.5% dose levels. The mean liver weights (absolute and relative to body weight) were different from the control mean (increased) at all dose levels, but no microscopic pathology was detected. Thyroid hyperplasia was reported in some animals at all doses, as was "very slight numerical development of the follicles and ripening follicles in the ovaries of females" at the high dose (4820 mg/kg/d). No gross or microscopic changes were detected in any other organ, and no change was detected in clinical chemistry tests.

The report concluded that "The increased liver weight must be attributed to hyperactivity; hypermetabolism as a result of increased thyroid activity appears probable in view of the observations of the thyroid". Therefore, the increased liver weights were not pathologic: there

were no microscopic lesions detected on histopathology and no change in clinical chemistry values. Recent work on the relationship of liver weight, microsomal enzyme induction, and histological change in rat toxicology studies has been published (Amacher et al, Food and Chemical Toxicology, 36, 831-839, 1998). This paper concluded "The preponderance of data collected in these 11 studies indicates that microsomal enzyme induction was not accompanied by evidence of chemically-induced liver injury. We conclude that in the rat, both hepatomegaly and microsomal enzyme induction are benign and adaptive changes in response to certain chemicals that stimulate the hepatic drug metabolizing enzyme system."

Conclusions

Input optional information or further explain the contents of a particular section, much as is done in the "Discussion" portion of a publication in academic journals.

The NOAEL in this 28-day study was 1% "Hexabromid S" in the diet. Based on body weights and food consumption data this dose is equivalent to 940 mg/kg body weight/day.

Data Quality

Reliability Reasonable

Denote the reliability of data, at the discretion of the person preparing the robust su

Data Reliability Remarks Add comments about how reliability of data was determined, or add re

This study was performed by a laboratory with considerable experience. However, the study was performed approximately 30 years ago using an HBCD product no longer manufactured as test article, and is not up to today's standards. The fact that recently conducted repeated dose studies with HBCD provided comparable results lends credence to the results of this study.

Reference

Cite the full reference for the critical study on which the robust study summary is based. List other appropriate references that support this summary and, have been reviewed for

>> Remarks

Zeller H and Kirsch P (1969) Hexabromocyclododecane: 28-day feeding trials with rats. BASF (unpublished laboratory report).

General

Add any statement that doesn't fit into any of the other f

This study was sponsored and performed by BASF.

The doses and results of this study are improperly reported by the Swedish Chemicals Inspectorate KEMI in the 1999 draft EU risk assessment of HBCD and in reports to the OECD SIDS programme. KEMI reports the doses as 0, 500, 1250 and 2500 mg/kg body weight/day (basis for conversion not given), and that the low-adverse-effect-level was 500 mg/kg (the 1% in the diet dose).

Sponsor ID	110401	Albemarle Corporation	Create Date	4/6/01
CAS Number	319155-5	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	4
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	

Revision Date:

12/6/01

Test Substance

Remarks

The test article was a commercial HBCD product ("Hexabromid S") produced at one time by BASF in Germany. BASF no longer manufactures HBCD.

Chemical Category

Method

>> Method/Guideline followed

Not known.

>> GLP Unknown

>> Year study performed 1970

Were Good Laboratory Practices followed in the st

>> Species

rat

>> Strain Mammal strain Sprague-Dawley

>> Sex

>> Number of males per dose >> Number of females per dose

>> Route of Administration

>> Exposure Period

Duration of study in days (for example, 28 days, 90 d)

>> Frequency of treatment

Number of doses per day, week, etc. This is particularly relevant for inhalation experiments -- 6hrs/day, 5 d

>>Doses

List all doses used in te

>> Control Group

Concurrent contro

>> Post observation period

Length of time animals observed after last d

>> Statistical Method

Cite statistical methods use

Remarks for Method

HBCD ("Hexabromid S") was tested in Sprague-Dawley rats at doses of 0, 0.16, 0.32, 0.64 and 1.28% of the diet for 90 days. Doses calculated on the actual body weights and food consumption in this study reveals: 0, 120, 240, 470 and 950 mg/kg body weight/day.

Results

>> NOAEL Precision

>> NOAEL dose >> Unit

>> NOAEL Effec

(e.g., decrease in body weight, organ

Increase in liver weight in the absence of pathology or clinical chemistry changes.

>> LOAEL Precision

>

>> LOAEL dose

1280

>> Unit

mg/kg in feed

>> LOAEL Effect

(e.g., decrease in body weight, organ

See Remarks.

>> Actual dose received by dose level by sex (if available)

0, 120, 240, 470 and 950 mg/kg body weight/day

>> Toxic response A brief narrative describing toxic response or effects, by dos

See Remarks.

>> Statistical results Note statistical results, with appropri ρ value

See Results Remarks.

Results Remark Provide at a minimum qualitative descriptions of elements where dose effect related observations of elements were seen:

HBCD ("Hexabromid S") was tested in Sprague-Dawley rats at doses of 0, 0.16, 0.32, 0.64 and 1.28% of the diet for 90 days. Doses calculated on the actual body weights and food consumption in this study reveals: 0, 120, 240, 470 and 950 mg/kg body weight/day.

Doses up to 0.64% (470 mg/kg/d) produced no adverse clinical signs, no change in body weight, and no change in clinical chemistry results. An increase in the relative liver to body weight ratio was found, and was accompanied by fatty accumulation but no other histologically discernible changes were detected in the liver. The pathology report states that although fat ("lipid phanerosis") was visible microscopically in the liver of treated rats, this change was not accompanied by any pathology and could not be defined as "fatty liver". Further, no histological changes were found in any other organ. The report states that in the "absence of detectable clinico-chemical disturbances or histological changes of the vital organs, it was concluded that

the increased liver weight and the fat deposits, both of which were largely reversible when administration of Hexabromid S was stopped, were the result of a temporary increase in the activity of the liver."

Conclusions

Input optional information or further explain the contents of a particular section, much as is done in the "Discussion" portion of a publication in academic journals.

The highest dose tested in the BASF 90 d study, 1.28% of the diet (950 mg/kg body weight/day), is the no adverse effect level or NOAEL.

Data Quality

Reliability Reasonable

Denote the reliability of data, at the discretion of the person preparing the robust su

Data Reliability Remarks Add comments about how reliability of data was determined, or add re

This study was performed by a laboratory with considerable experience. However, the study was performed approximately 30 years ago using an HBCD product no longer manufactured as test article, and is not up to today's standards. The fact that recently conducted repeated dose studies with HBCD provided comparable results lends credence to the results of this study.

Reference

Cite the full reference for the critical study on which the robust study summary is based. List other appropriate references that support this summary and, have been reviewed for

>> Remarks

Zeller H and Kirsch P (1970) Hexabromocyclododecane: 90-day feeding trials with rats. BASF (unpublished laboratory report).

General

Add any statement that doesn't fit into any of the other f

This study was sponsored and performed by BASF.

The Swedish Chemicals Inspectorate (KEMI) improperly reported the results of this study and incorrectly converted the doses from a percentage of the diet to mg/kg body weight in 1999 draft EU risk assessment on HBCD and in reports to the OECD SIDS programme. KEMI converted the dietary doses to 0, 80, 160, 320 and 640 mg/kg body weight/day (basis for conversion not given). KEMI also stated that the low-adverse-effect level (LOAEL) in this

study was 80 mg/kg (the 0.16% in the diet dose level, and that a no effect level (NOEL) was not determined in any of the subchronic studies conducted to date, including the 1997 Rat 28-Day Study.

The results of the BASF 90 day study do not indicate that an adverse effect was produced at 0.16% of the diet. Further, the results indicate no adverse effect was produced at the highest dose tested, 1.28% of the diet. The BASF pathology report clearly states that although fat ("lipid phanerosis") was visible microscopically in the liver of treated rats, this change was not accompanied by any pathology and could not be defined as "fatty liver". Therefore, not even the highest dose tested in the 90-day study can be defined as the low adverse effect level (LOAEL). The highest dose tested in the BASF 90 d study, 1.28%, is more accurately defined as the no adverse effect level or NOAEL.

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1106021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

EPA OPPTS Method 870.5375 In vitro Mammalian Chromosome Aberration Test

>> Test Type

Cytogenetic assay

>> System of Testing

Non-bacterial

>> GLP

Yes

>> Year study performed

1996

>> Species

Primary cultures - human lymphocytes

>> Metabolic Activation

Arochlor 1254-induced rat liver S-9; prepared from male Sprague-Dawley rats

>> Concentration

Initial: 75, 250, 750, 2500 ug/ml; Definitive: 10, 19, 38, 75, 150, 300, 600 ug/ml

>> Statistical Method

Fisher's exact test

Remarks for Method

The test article, Hexabromocyclododecane (HBCD) was tested in the in vitro mammalian cytogenetic test using human peripheral blood lymphocytes (HPBL) in both the absence and presence of metabolic activation. The assay was performed in two phases. The first phase, the initial chromosome aberration assay, was conducted to establish the dose range for testing and to evaluate the clastogenic potential of the test article. The second phase, the independent repeat chromosome aberration assay, was performed to confirm the test system response to the test article seen in the initial assay.

12/20/01

Page 1 of 1

0

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity in Vitro (Gene Mutations)

Sponsor ID	1103021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Dimethylsulfoxide (DMSO) was the solvent of choice based on the solubility of the test article and compatability with the target cells. The test article was soluble in DMSO at ~500 mg/ml, the highest concentration tested.

Initial. In the initial chromosome aberration assay, duplicate cultures of HPBL were exposed to 9 concentrations of the test article, and to positive, solvent and negative controls. The dividing cells were harvested at ~20 hours after initiation of treatment. The maximum dose tested was 2500 ug/ml. Dose levels greater than 2500 ug/ml were insoluble in the treatment medium and not tested. Visible precipitate was observed in treatment medium at dose levels of 750 and 2500 ug/ml and was soluble but cloudy (no visible precipitate) at dose levels 75 and 250 ug/ml. The test article was soluble in treatment medium at all other dose levels tested. In the non-activated portion of the test, HPBL cells were exposed to the test article continuously for 20 hours; in the S9-activated portion of the test, HPBL were exposed to the test article for 4 hours. Metaphase cells were collected for microscopic evaluation at 20 hours after the initiation of treatment.

Second Phase. Duplicate cultures of HPBL were exposed to at least 4 concentrations of the test article, as well as solvent, positive, and untreated controls. The dose levels selected were based on the initial assay. The dividing cells were harvested at 2 time points: 20 and 44 hours after initiation of treatment. HBCD was tested in the absence and presence of an Arochlor-induced S9 metabolic activation system at dose levels of 10, 19, 38, 75, 150, 300 and 600 ug/ml. The test article was soluble but cloudy at 75 ug/ml and was workable in treatment medium at dose levels 150 ug/ml and higher. The test article was soluble in treatment medium at all other concentrations tested. In the independent repeat assay, HPBL cells were exposed to the test article continuously for 20 or 44 hours in the non-activated test system and for 4 hours in the S9-activated test system. Metaphase cells were collected for microscopic evaluation in both the non-activated and S9-activated studies at 20 and 44 hours after the initiation of treatment.

Evaluation of Metaphase Cells. Metaphase cells with 46 centromeres were examined under oil immersion without knowledge of treatment groups. Whenever possible, a minimum of 200 metaphase spreads (100 per duplicate treatment condition) were examined and scored for chromatid-type and chromosome-type aberrations. The mitotic index was recorded as the percentage of cells in mitosis per 500 cells counted. In the delayed harvests, the percent polyploid cells was recorded per 100 metaphase cells.

Controls. Mitomycin C was used as the positive control in the non-activated study. Cyclophosphamide was used as the positive control in the S-9 activated study. For both positive controls one dose with sufficient scorable metaphase cells was selected for analysis. The solvent vehicle for the test article was used as the solvent control at the same concentration as that found in the test article-treated groups. Growth medium or S9 reaction mixture was used in the untreated control.

Evaluation of Results. Toxic effects of treatment were based on mitotic inhibition relative to the solvent-treated control. The number and types of aberrations, the percent aberrant cells, the percentage of numerically damaged cells and the frequency of structural aberrations per cell

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	319356	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

was reported for each treatment group.

Results

>> Result Negative

>> Cytotoxic Concentration

Non-activated: toxicity at 750 ug/ml; S9-activated: toxicity at 250 ug/ml

>> Genotoxic Effects Unconfirmed

>> Statistical results

No statistically significant differences were observed between the negative, solvent and treatment groups ($p > 0.05$, Fisher's exact test). The positive controls performed as expected.

Results Remark

In the initial assay, dose levels of 2500 ug/ml in the non-activated study and 750 and 2500 ug/ml in the S9-activated study were not analyzed from chromosome aberrations due to complete mitotic inhibition. Toxicity (mitotic inhibition) of ~56% was observed at the highest dose level (750 ug/ml) evaluated for chromosome aberrations, in the non-activated study. In the S9-activated study, 13% toxicity was observed at the highest dose level (250 ug/ml) evaluated for chromosome aberrations. No statistically significant increases in chromosome aberrations were observed in either the non-activated or S9-activated test systems relative to the solvent control group regardless of dose level ($p > 0.05$, Fisher's exact test).

In the independent repeat chromosome aberration assay, toxicity, as measured by mitotic inhibition, was ~55% and 94% at the 20 and 44 hour harvest, respectively, at the highest dose levels (600 and 300 ug/ml) evaluated in the non-activated studies. In the S9-activated studies, toxicity was approximately 71% and 69% at the 20 and 44 hour harvest, respectively, at the highest dose levels (300 and 600 ug/ml) evaluated. The 600 ug/ml dose level in the non-activated 44 hour harvest and in the S9-activated 20 hour harvest was not analyzed for chromosome aberrations due to an insufficient number of scorable metaphase cells. No statistically significant increases in structural chromosome aberrations were observed in either the non-activated or S9-activated studies, regardless of dose level or harvest time ($p > 0.05$, Fisher's exact test). No statistically significant increases in numerical chromosome aberrations were observed in either the non-activated or S9-activated studies at the 44 hour harvest time, regardless of dose level ($p > 0.05$, Fisher's exact test).

Conclusions

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

HBCD was negative for the induction of structural and numerical chromosome aberrations in human peripheral blood lymphocytes.

Data Quality

Reliability High

Data Reliability Remarks

This study was performed using current techniques, under Good Laboratory Practices, by a laboratory with considerable experience performing this type of study.

Reference

>> Remarks

Gudi, R. And Schady, E. (1996) Chromosome Aberrations in Human Peripheral Blood Lymphocytes. Hexabromocyclododecane. Laboratory Study Number G96AO61.342. Microbiological Associates, Inc., Rockville, MD.

General

Study sponsored by the Chemical Manufacturers Association Brominated Flame Retardant Industry Panel.

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	11/01
CAS Number	119-1556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

12/5/01

Test Substance

Remarks

Exact composition of the test article is not known.

Chemical Category

Method

>> Method/Guideline followed

Not specified

>> Test Type

Ames test

>> System of Testing Bacterial

>> GLP Unknown

>> Year study performed 1976

>> Species

Salmonella typhimurium

>> Metabolic Activation

Arochlor induced rat liver S9

>> Concentration

0, 1, 10, 50, 100, 500, 1000, 5000 ug/plate

>> Statistical Method

Not known.

Remarks for Method

Five strains of Salmonella typhimurium (TA1535, TA1537, TA1538, TA98 and TA100) were tested in the presence and absence of a metabolic activation system (Arochlor induced rat liver). Doses were 0, 1, 10, 50, 100, 500, 1000 or 5000 ug HBCD/plate.

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3193556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Results

>> Result Negative

>> Cytotoxic Concentration

>5000 ug HBCD/plate with or without metabolic activation

>> Genotoxic Effects Unconfirmed

>> Statistical results

Not known.

Results Remark

The test article was not mutagenic or toxic at any dose level when tested with or without metabolic activation.

Conclusions

HBCD was not mutagenic in *S. typhimurium* at doses up to and including 5000 ug/plate when tested with or without metabolic activation. These results are consistent with other Ames test's performed on this material (Ogaswara S and Hanafusa T. (1993) Report on mutagenicity test on Pyroguard SR-103 using microorganisms; Baskin A and Phillips, B. (1977) Mutagenicity of two lots of FM-100, Lot 53 and residue of Lot 3322 in the absence and presence of metabolic activation. Industrial Biotest Laboratories, Sponsored by Velsicol Chemical Corporation; Anonymous. (1979) Mutagenicity test of GLS-S6-41A. Gulf South Research Institute, Sponsored by Ethyl Corporation; US Environmental Protection Agency (1990) Ames metabolic activation test to assess the potential mutagenic effect of Compound No. 49. Letter from BASF. EPA/OTS Doc #86-900000385.

Data Quality

Reliability

Acceptable

Data Reliability Remarks

Multiple Ames tests performed at different test laboratories using different commercial HBCD products as test article have all been negative. The consistency of the negative results increases the confidence in the results.

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	1194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	2
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Reference

>> Remarks

Simmons V., Poole, D., Newell, G., and Skinner, W. (1976) In vitro microbiological mutagenicity studies for four CIBA-GEIGY Corporation compounds. SRI Project LSC-5702.

General

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	3191556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

12/6/01

Test Substance

Remarks

HBCD, obtained from Aldrich Chemicals (Stockholm, Sweden)

Chemical Category

Method

>> Method/Guideline followed

Non standard test methodology

>> Test Type

Mammalian cells in culture (Sp5 and SPD8 duplication cell lines)

>> System of Testing

Non-bacterial

>> GLP

No

>> Year study performed

1999

>> Species

Not known.

>> Metabolic Activation

None

>> Concentration

See results.

>> Statistical Method

Student's t test

Remarks for Method

HBCD was tested in vitro in hamster cells (Sp5/V79 and SPD8) in a recombination assay at five doses between 2 and 20 ug/ml plus a control. The Sp5 and SPD8 clones exhibit a spontaneous partial duplication of the HPRT gene, resulting in a non-functional HGPRT protein. The mutants revert spontaneously to a functional HPRT gene phenotype by recombination; an increase in reversion frequency is considered a positive response.

Treatment with HBCD resulted in a ~ maximal 2-fold increase in revertant frequency, which

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1100921	Albemarle Corporation	Create Date	1/6/01
CAS Number	319156	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	3
Consortia ID	1101912	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

was reported as statistically significant.

This reliability of this genetic test is unknown. The reproducibility of the results, validation of the system, dose-effect response, and whether a maximal two-fold increase is evidence of a positive response are also unknown.

Results

>> Result Ambiguous

>> Cytotoxic Concentration

Not known.

>> Genotoxic Effects Equivocal

>> Statistical results

See Remarks Section.

Results Remark

Treatment with HBCD resulted in a ~ maximal 2-fold increase in revertant frequency, which was reported as statistically significant.

This reliability of this genetic test is unknown. The reproducibility of the results, validation of the system, dose-effect response, and whether a maximal two-fold increase is evidence of a positive response are also unknown.

Conclusions

Treatment with HBCD resulted in a ~ maximal 2-fold increase in revertant frequency, which was reported as statistically significant. The reliability of this test is unknown.

Data Quality

Reliability Unknown.

Data Reliability Remarks

EPA High Production Volume (HPV) Track

Toxicity End point:
Toxicity In Vitro (Gene Mutations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	4/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	3
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed	Y

The reliability and predictive ability of this genetic test is unknown. The reproducibility of the results, validation of the system, dose-effect response, and whether a maximal two-fold increase is evidence of a positive response are unknown.

Reference

>> Remarks

Helleday et al, Mutat Res, 1999, 439(2): 137-147.

General

EPA High Production Volume (HPV) Track

Toxicity End Point:
Toxicity In Vivo (Chromosomal Aberrations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	11/6/01
CAS Number	3194556	Cyclododecane, 1,2,5,6,9,10-hexabromo	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Revision Date:

12/5/01

Test Substance

Remarks

The test article was a composite of equal parts of the commercial hexabromocyclododecane (HBCD) commercial product produced by Albemarle Corporation, Dead Sea Bromine Group, and Great Lakes Chemical Corporation. The test article composite was analyzed for characterization and homogeneity. The results of the analysis indicated the test article was homogeneous and contained the following components: HBCD beta isomer 8.5%, HBCD alpha isomer 6.0%, HBCD gamma isomer 79.1%.

Chemical Category

Method

>> Method/Guideline followed

OECD Method 474

>> Test Type

Micronucleus assay

>> GLP Yes

>> Year study performed 2000

>> Species

mouse

>> Strain Mammal strai NMRI

>> Sex M

>> Number of males per dose

5

>> Number of females per dose

0

>> Route of Administration

Intraperitoneal

>> Doses 0, 500, 1000, 2000 mg/kg

>> Exposure period

Two doses administered 24 hrs apart.

>> Statistical Method

Wilcoxon test

Remarks for Method

EPA High Production Volume (HPV) Track

Toxicity End Point:
Toxicity in Vivo (Chromosomal Aberrations)

Sponsor ID	1100023	Albemarle Corporation	Create Date	1/6/01
CAS Number	3193576	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Hexabromocyclododecane (HBCD) was tested for clastogenicity and for the ability to induce spindle poison effects in NMRI mice (Charles River Deutschland GmbH) using the micronucleus method. HBCD, dissolved in DMSO, was administered twice intraperitoneally with a 24-hr interval between doses to male mice (n=5/group) at dose levels of 500, 1000 or 2000 mg/kg body weight in a volume of 4 ml/kg. DMSO (the vehicle) was administered to male mice by the same route and frequency. Cyclophosphamide was used as a positive control for clastogenic effects. Vincristine was used as a positive control for induction of spindle poison effects. Animals in the positive control groups were treated only once.

The animals were sacrificed and the bone marrow of the two femora prepared 24 hours after the second administration. After staining, 2000 polychromatic erythrocytes were evaluated per animal and investigated for micronuclei. The normocytes with and without micronuclei occurring per 2000 polychromatic erythrocytes were also counted.

Results

>> Effects on Mitosi

PCE/NCE 0, 500, 1000, 2000 mg/kg = 3.74, 2.89, 2.67, 2.49, respectively.

>> Genotoxic Effects

Negative

>> Statistical results

No statistical differences between the treatment and vehicle control group were observed ($p \leq 0.05$).

Results Remark

The two intraperitoneal administrations of DMSO in a volume of 4 ml/kg body weight led to 1.4% polychromatic erythrocytes containing micronuclei. In the 2000 mg HBCD/kg body weight group, 0.9% micronuclei were found. In the 1000 and 500 mg HBCD/kg body weight groups, 1.0 and 1.1% micronuclei were detected. The two positive control substances performed as expected.

The number of normochromatic erythrocytes containing micronuclei did not differ to any appreciable extent in the negative control or various dose groups.

Conclusions

HBCD treatment did not increase numbers of micronuclei. The number of normochromatic or polychromatic erythrocytes containing small micronuclei did not deviate from the vehicle control value and was within the historical control range. Large micronuclei were not observed. HBCD had no chromosome-damaging (clastogenic) effect in this study and did not impair chromosome distribution during mitosis.

EPA High Production Volume (HPV) Track

Toxicity End Point:
Toxicity In Vivo (Chromosomal Aberrations)

Sponsor ID	1100021	Albemarle Corporation	Create Date	1/6/01
CAS Number	319-1556	Cyclododecane, 1,2,5,6,9,10-hexabromo-	Study Number	1
Consortia ID	1101012	CMA Brominated Flame Retardant Industry Panel (BFRIP)	Completed:	Y

Data Quality

Reliability High

Data Reliability Remarks

This study was performed according to current guidelines under Good Laboratory Practices by an experienced laboratory.

Reference

>> Remarks

Engelhardt, G and Hoffmann, H. (2000) Cytogenetic Study in vivo with Hexabromocyclododecane in the Mouse Micronucleus Test After Two Intraperitoneal Administrations. Laboratory Project Identification: 26M0100/004018. Experimental Toxicology and Ecology, BASF Aktiengesellschaft, Ludwigshafen, Germany.

General